

# Contact of a regular surface in Euclidean 3-space with cylinders and cubic binary differential equations

T. Fukui\*, M. Hasegawa†, and K. Nakagawa

November 28, 2014

## Abstract

We investigate contact type of a regular surface in Euclidean 3-space  $\mathbb{R}^3$  with right circular cylinders. We present the conditions for existence of cylinders with  $A_1$ ,  $A_2$ ,  $A_3$ ,  $A_4$ ,  $A_5$ ,  $D_4$ , and  $D_5$  contacts with given surface. We also investigate the kernel field of  $A_{\geq 3}$ -contact cylinders on the surface. This is defined by a cubic binary differential equation and we classify singularity types of its flow in generic context.

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Monge normal form</b>	<b>3</b>
<b>3</b>	<b>Contact</b>	<b>4</b>
3.1	Definition of contact type . . . . .	4
3.2	Contact with spheres . . . . .	4
3.3	Contact with planes . . . . .	5
<b>4</b>	<b>Contact with cylinders</b>	<b>5</b>
4.1	Main results . . . . .	6
4.2	Contact map with cylinders and its singularity type . . . . .	8
4.2.1	Case (a): $k_1 k_2 \neq 0$ , $\lambda = \frac{\kappa_n(\mathbf{w})}{k_1 k_2}$ . . . . .	10
4.2.2	Case (b): $k_1 = 0$ , $k_2 \neq 0$ , $\mathbf{w} = \partial_u$ . . . . .	11
4.2.3	Case (c): $k_1 = k_2 = 0$ . . . . .	13

---

2010 *Mathematics subject classification.* 53A05,53A60,58K05,58C27

*Key words and phrases.* contact with cylinders, Monge cubic, cylindrical directions.

\*Partially supported by Grant-in-Aid in Sciences 24540067.

†Supported by the FAPESP post-doctoral grant number 2013/02543-1. The post-doctoral is held at ICM- USP, Sao Carlos, Brazil.

<b>5</b>	<b>Cylindrical directions</b>	<b>14</b>
5.1	Monge normal form of a surface given by Monge normal form . . . . .	14
5.2	Monge cubic . . . . .	19
5.3	Conjugate Monge cubic . . . . .	21
5.4	Reduction of jets of cubic differential equations . . . . .	25

# 1 Introduction

In his celebrated book [19, page 96], R. Thom pointed out the similarities of the geometric features of  $D_4$  singularities and umbilic of regular surface in  $\mathbb{R}^3$ . Inspired this observation, I. Porteous ([16, 17]) showed that many differential geometric concepts are rediscovered by looking the singularities of distance-squared functions. He found new notion called by ridge which corresponds to  $A_3$ -singularities. After J. Montaldi ([15]) defines the notion of contact between two submanifolds and established the relation to  $\mathcal{K}$ -equivalence, introduced by J. Mather ([14, §2]), in singularity theory, this is justified as an investigation of contact of a surface with spheres. The contact with plane is also recognized fundamental and the second fundamental form plays crucial rule. See [3], for example, for several relations of singularities of height function and differential geometric concepts,

A homogeneous surface in Euclidean 3-space  $\mathbb{R}^3$  is an orbit of a certain subgroup of Euclidean motion group  $O(3) \times \mathbb{R}^3$ . They are planes, spheres or cylinders ([18]).

Investigating contacts of a surface with planes and spheres has produced a rich field connecting differential geometry with singularity theory and this has been done by many authors (see [9, 10] for typical research articles, [2] for a standard textbook, [7, 8] for surveys, and their references). In this paper, we will investigate the contact of a surface with cylinders, which has not been investigated in detail yet.

Our computation is based on Monge normal form and this is given in §2 with definitions of several concepts. In §3, we define the notion of contact due to J. Montaldi and quickly recall the conditions that a surface has a sphere or plane with degenerate contact in generic context. Then we state our main theorem in §4.1. The proof of main theorem will be given in §4.2.

On a surface in  $\mathbb{R}^3$ , there are many articles discussing direction fields defined by quadratic differential equations (see e.g. [4] and its references), like principal directions, asymptotic directions and characteristic directions. As far as we know, there are no literatures concerning a natural direction fields on a surface in  $\mathbb{R}^3$  defined by a cubic differential equation in a geometric context. The kernel field of  $A_2$ -contact with spheres defines principal directions when the point is not umbilic. The kernel field of  $A_1^-$ -contact with planes defines asymptotic directions in the hyperbolic region. So it is natural to investigate kernel fields of  $A_3$ -contact with cylinders. We call them by **cylindrical fields**, and we show that they are defined by a cubic differential equation.

The last section is dedicated to analysis of the singularities of integral curves of

cylindrical fields in generic context. We show that singularities of flows of cylindrical fields are on the discriminant of Monge cubic or parabolic lines. To do this, we present a formula for Monge normal form at a non-umbilical point in a surface given by Monge normal form in §5.1. We discuss several explicit method of determination of singularity types of the flow.

## 2 Monge normal form

Let  $\langle \cdot, \cdot \rangle$  denote Euclidean inner product. For any regular surface  $S$  in Euclidean 3-space  $\mathbb{R}^3$  and for any point  $\mathbf{p}$  on  $S$ , we can find an isometric coordinate change  $T : (\mathbb{R}^3, \mathbf{p}) \rightarrow (\mathbb{R}^3, 0)$  so that  $T(S)$  is expressed by the image of the map

$$\mathbf{x} : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^3, 0), (u, v) \mapsto (u, v, f(u, v)), \quad (2.1)$$

at least locally, where  $f(u, v) = \frac{1}{2}(k_1u^2 + k_2v^2) + \sum_{s=3}^k c_s + o(k)$ ,  $c_s = \sum_{i+j=s} \frac{a_{i,j}}{i!j!} u^i v^j$ . The expression (2.1) is called by Monge normal form and the coefficients  $k_1, k_2, a_{i,j}$  are differential-geometric invariants of  $S$  at  $\mathbf{p}$ , at least when one assume that  $k_1 < k_2$ . Remark that  $k_1$  and  $k_2$  are principal curvatures at  $(0, 0)$ . We set  $\mathbf{v}_1 = \partial_u$  and  $\mathbf{v}_2 = \partial_v$ . These are principal vectors at  $(0, 0)$  with respect to  $k_1$  and  $k_2$  respectively.

For a regular surface  $S$  in  $\mathbb{R}^3$  defined by (2.1), the second fundamental form of this surface at  $(0, 0)$  is the bilinear form defined by

$$(\mathbf{w}, \mathbf{w}') \mapsto \text{II}(\mathbf{w}, \mathbf{w}') = k_1 w_1 w_1' + k_2 w_2 w_2'$$

where  $\mathbf{w} = w_1 \partial_u + w_2 \partial_v$ ,  $\mathbf{w}' = w_1' \partial_u + w_2' \partial_v$ . We say  $\mathbf{w}$  is asymptotic if  $\text{II}(\mathbf{w}, \mathbf{w}) = 0$ .

We say  $(0, 0)$  is **ridge** with respect to  $\mathbf{v}_1$  (resp.  $\mathbf{v}_2$ ), if  $a_{30} = 0$  (resp.  $a_{03} = 0$ ). This condition is equivalent that  $\mathbf{v}_1 \kappa_1 = 0$  (resp.  $\mathbf{v}_2 \kappa_2 = 0$ ) at  $(0, 0)$  where  $\kappa_i = \kappa_i(u, v)$  is the principal curvature with  $\kappa_i(0, 0) = k_i$ . We say the ridge is of the first order if  $\mathbf{v}_i \kappa_i(0, 0) = 0$ ,  $\mathbf{v}_i^2 \kappa_i(0, 0) \neq 0$ . More generally, we say that  $(0, 0)$  is the  $k$ -th order ridge with respect to  $\mathbf{v}_i$  if  $\mathbf{v}_i^j \kappa_i(0, 0) = 0$  if  $j \leq k$  and  $\mathbf{v}_i^{k+1} \kappa_i(0, 0) \neq 0$ .

We say  $(0, 0)$  is **subparabolic** with respect to  $\mathbf{v}_j$  if  $\mathbf{v}_j \kappa_i(0, 0) = 0$  ( $\{i, j\} = \{1, 2\}$ ). This is equivalent that  $a_{ij} = 0$ .

When we assume that  $k_1 k_2 \neq 0$ , we can define the conjugate direction. The **conjugate direction** of the direction generated by a vector  $\mathbf{w}$  is the direction generated by a non-zero vector  $\bar{\mathbf{w}}$  with  $\text{II}(\mathbf{w}, \bar{\mathbf{w}}) = 0$ . So if  $\mathbf{w} = w_1 \partial_u + w_2 \partial_v$ , then the conjugate direction is generated by  $k_2 w_2 \partial_u - k_1 w_1 \partial_v$ .

We often refer  $6c_3$  as **Monge cubic** of  $S$  at  $(0, 0)$ . We call  $\bar{c}_3(\mathbf{w}) = 6c_3(\bar{\mathbf{w}})$  the **conjugate Monge cubic** at  $(0, 0)$ , and the root(s) of this cubic  $\bar{c}_3$  **cylindrical direction(s)** (or, **generatrix direction(s)**).

For a function  $\varphi(u, v)$ , we set  $\nabla \varphi(u, v) = \varphi_u(u, v) \partial_u + \varphi_v(u, v) \partial_v$ .

### 3 Contact

#### 3.1 Definition of contact type


According to [15], we define the notion of contact type. Given two pairs of submanifold germs at the origin in  $\mathbb{R}^n$ , the pairs have the same contact type if there is a diffeomorphism-germ of  $(\mathbb{R}^n, 0)$  taking one pair to the other.

**Theorem 3.1** ([15]). *For  $i = 1, 2$ , let  $g_i : (X_i, x_i) \rightarrow (\mathbb{R}^n, 0)$  be immersion-germs and  $f_i : (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^p, 0)$  be submersion-germs with  $Y_i = f_i^{-1}(0)$ . Then the pairs  $(X_1, Y_1)$  and  $(X_2, Y_2)$  have the same contact type if and only if  $f_1 \circ g_1$  and  $f_2 \circ g_2$  are  $\mathcal{K}$ -equivalent.*

Here we say that two map-germs  $f, g : (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^p, 0)$  are  **$\mathcal{K}$ -equivalent** if there are a diffeomorphism  $\phi : (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^n, 0)$  and a smooth map  $A : (\mathbb{R}^n, 0) \rightarrow \text{GL}(\mathbb{R}^p)$  so that  $g(\phi(x)) = A(x)f(x)$ . In this paper, we consider the following contact types corresponding to  $\mathcal{K}$ -equivalence classes of  $(\mathbb{R}^2, 0) \rightarrow (\mathbb{R}, 0)$  represented by

$$A_k \text{ (or } A_k^\pm) : x^2 \pm y^{k+1}, \quad D_k \text{ (or } D_k^\pm) : x(y^2 \pm x^{k-2}), (k \geq 4).$$

We show below the zero set of these singularities. Remark that  $A_k$  singularities define kernel field of Hessian when  $k \geq 2$ .



#### 3.2 Contact with spheres

Since spheres in Euclidean 3-space are determined by their centers and radii the moduli space of spheres is of four dimensional. So we expect that there are  $A_1, A_2, A_3, A_4, D_4$ -contact spheres for a generic surface.


In [16, 17], I. Porteous had investigated the singularities of distance squared functions on the surface and these functions measure the contact with sphere. We quickly recall some of his results here.

Let  $S_{a,b,c}$  denote the sphere centred at  $(a, b, c)$  with radius  $\sqrt{a^2 + b^2 + c^2}$ . For a regular surface  $S$  in  $\mathbb{R}^3$  defined by (2.1), we have the following.

1. The sphere  $S_{a,b,c}$  is  $A_{\geq 1}$ -contact with  $S$  at  $(0, 0, 0)$  when  $(a, b, c)$  is on the normal line.
2. The sphere  $S_{a,b,c}$  is  $A_{\geq 2}$ -contact with  $S$  at  $(0, 0, 0)$  when  $(a, b, c)$  is on the focal set of  $S$  on the normal line, (i.e., the distance between  $(a, b, c)$  and  $(0, 0, 0)$  is  $1/k_i$ ). This sphere is often referred as a curvature sphere with respect to  $k_i$ .
3. If  $(0, 0)$  is ridge with respect to  $\mathbf{v}_i$  (i.e.,  $\mathbf{v}_i \kappa_i(0, 0) = 0$ ), the surface  $S$  is  $A_{\geq 3}$ -contact with the curvature sphere  $S_{a,b,c}$  with respect to  $k_i$  at  $(0, 0, 0)$ .

4. If  $(0,0)$  is at least second order ridge with respect to  $\mathbf{v}_i$  (i.e.,  $\mathbf{v}_i \kappa_i(0,0) = \mathbf{v}_i^2 \kappa_i(0,0) = 0$ ), the surface  $S$  is  $A_{\geq 4}$ -contact with the curvature sphere  $S_{a,b,c}$  with respect to  $k_i$  at  $(0,0,0)$ .
5. If  $(0,0)$  is an umbilic, the curvature sphere  $S_{a,b,c}$  has  $D_{\geq 4}$ -contact with  $S$  at  $(0,0,0)$ .

Remark that the kernel field of  $A_{\geq 2}$ -contact with sphere is defined except umbilics and it defines principal fields. Classification of singularity types of the curvature curves at generic umbilics are known as Darboux classification ([5, Note VII]).



### 3.3 Contact with planes

Since planes in Euclidean 3-space are defined by  $ax + by + cz = d$ , the moduli space of planes is of three dimensional. So we expect that there are  $A_1, A_2, A_3$ -contact planes for a generic surface.


To investigate height functions on the surface, we measure contact with planes. We also recall it quickly. Let  $\pi_{a,b,c}$  denote the plane defined by  $ax + by + cz = 0$  with  $a^2 + b^2 + c^2 = 1$ .

For a regular surface  $S$  in  $\mathbb{R}^3$  defined by (2.1), we have the following.

1. The plane  $\pi_{a,b,c}$  is  $A_{\geq 1}$ -contact with  $S$  at  $(0,0,0)$  when  $(a,b,c)$  is on the normal line.
2. The plane  $\pi_{a,b,c}$  is  $A_{\geq 2}$ -contact with  $S$  at  $(0,0,0)$  when  $(0,0,0)$  is parabolic.
3. If  $(0,0)$  is parabolic (i.e.,  $k_i = 0$ ) and ridge with respect to  $\mathbf{v}_i$  (i.e.,  $\mathbf{v}_i \kappa_i(0,0) = 0$ ), the surface  $S$  is  $A_{\geq 3}$ -contact with the plane  $\pi_{a,b,c}$  at  $(0,0,0)$ .

Remark that the kernel field of  $A_1^-$ -contact with plane is defined in hyperbolic region and it defines asymptotic directions.

Classification of singularity types of asymptotic curves are given by [1] as follows.



## 4 Contact with cylinders

Now we consider the contact type of a regular surface with cylinders. Since cylinders in Euclidean 3-space are determines by their central axes and radii, and central axes

are elements of Grassmannian, the moduli space of cylinders is of five dimensional. So we expect that there are  $A_1, A_2, A_3, A_4, A_5, D_4, D_5$ -contact cylinders for a generic surface.

It is clear that a regular surface  $S$  and a cylinder  $C$  meet transversely at a point  $P$  if they do not have the common normal line at  $P$ . So we consider the contact of  $S$  and  $C$  with common normal line at  $P \in S \cap C$ .

## 4.1 Main results

For a unit vector  $\mathbf{w} = w_1\partial_u + w_2\partial_v$ , we consider the orthogonal projection

$$P_{\mathbf{w}} : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^2, 0), (u, v) \mapsto (w_2u - w_1v, f(u, v)).$$

When  $(0, 0)$  is ridge with respect to a principal vector  $\mathbf{w} = \mathbf{v}_i$ . The critical locus of  $P_{\mathbf{w}}$  has a local branch whose tangent is the line generated by  $\mathbf{w}$ . Let  $D'_{\mathbf{w}}$  denote its image by  $P_{\mathbf{w}}$ .

The main results are stated as follows.

**Theorem 4.1.** *Let  $S$  be a regular surface defined by (2.1). Let  $C_{\mathbf{w}, \lambda}$  denote the cylinder which contains  $(0, 0, 0)$  and whose central axis is parallel to a unit vector  $\mathbf{w}$ , and contains a point  $(0, 0, \lambda)$ ,  $\lambda \neq 0$ .*

1. *The cylinder  $C_{\mathbf{w}, \lambda}$  has  $A_1$ -contact with  $S$  at  $(0, 0, 0)$  if and only if  $k_1k_2\lambda \neq \kappa_n(\mathbf{w})$  where  $\kappa_n(\mathbf{w}) = \Pi(\mathbf{w}, \mathbf{w})$ .*
2. *The cylinder  $C_{\mathbf{w}, \lambda}$  has  $A_2$ -contact with  $S$  at  $(0, 0, 0)$  if and only if one of the following conditions hold.*

*(A<sub>2a</sub>)  $(0, 0)$  is not parabolic (i.e.,  $k_1k_2 \neq 0$ ),  $\mathbf{w}$  is not cylindrical, and  $1/\lambda$  is the cylindrical curvature with respect to  $\mathbf{w}$  (i.e.,  $\lambda = \kappa_n(\mathbf{w})/(k_1k_2)$ ), see the definition at the beginning of subsection 4.2.1).*

*(A<sub>2b</sub>)  $(0, 0)$  is parabolic but not umbilic (i.e.,  $k_i = 0, k_j \neq 0, i \neq j$ ),  $w_j = 0, \lambda \neq 1/k_j$ , and  $(0, 0)$  is not ridge with respect to  $\mathbf{v}_i$ .*

*(A<sub>2c</sub>)  $(0, 0)$  is a flat umbilic (i.e.,  $k_1 = k_2 = 0$ ), and  $c_3(\mathbf{w}) \neq 0$ .*

3. *The cylinder  $C_{\mathbf{w}, \lambda}$  has  $A_3$ -contact with  $S$  at  $(0, 0, 0)$  if and only if one of the following conditions hold.*

*(A<sub>3a</sub>)  $(0, 0)$  is not parabolic (i.e.,  $k_1k_2 \neq 0$ ),  $\mathbf{w}$  is cylindrical,  $1/\lambda$  is cylindrical curvature with respect to  $\mathbf{w}$ , and the critical value set of the orthogonal projection with respect to  $\mathbf{w}$  is nonsingular and is the first order vertex (there is a circle with 4-point contact).*

*(A<sub>3b</sub>)  $(0, 0)$  is parabolic but not umbilic (i.e.,  $k_i = 0, k_j \neq 0, i \neq j$ ),  $\mathbf{w}$  is parallel to the principal vector  $\mathbf{v}_i$  (i.e.,  $w_j = 0$ ),  $1/\lambda$  is not a principal curvature (i.e.,  $1/\lambda \neq k_j$ ),  $(0, 0)$  is ridge with respect to  $\mathbf{v}_i$ , and*

- *the line generated by  $\mathbf{w}$  has 4-point contact with  $S$ , and  $1/\lambda$  is not the limit of curvature  $D'_{\mathbf{w}}$  at  $(0, 0)$ , or*

- the line generated by  $\mathbf{w}$  has at least 5-point contact with  $S$ , and  $(0,0)$  is not  $\mathbf{v}_j$ -subparabolic.

(A<sub>3c</sub>)  $(0,0)$  is a flat umbilic (i.e.,  $k_1 = k_2 = 0$ ),  $c_3(\mathbf{w}) = 0$ , and  $2c_4(\mathbf{w}) + \lambda \langle J\mathbf{w}, \nabla c_3(\mathbf{w}) \rangle^2 \neq 0$  where  $J\mathbf{w} = w_2\partial_u - w_1\partial_v$ .

**Remark 4.2.** It is worth to point out that there are no flat umbilics for general surfaces. This means that flat umbilics disappear under small perturbation. In other words, being a flat umbilic is not a generic conditions.

**Theorem 4.3.** The cylinder  $C_{\mathbf{w},\lambda}$  has  $D_4$ -contact with  $S$  at  $(0,0,0)$  if and only if  $(0,0)$  is parabolic but not umbilic (i.e.,  $k_i = 0, k_j \neq 0, i \neq j$ ),  $\mathbf{w}$  is parallel to the principal vector  $\mathbf{v}_i$  (i.e.,  $w_j = 0$ ),  $1/\lambda$  is the other principal curvature (i.e.,  $1/\lambda = k_j$ ), and  $c_3$  does not have multiple roots.

**Theorem 4.4.** The cylinder  $C_{\mathbf{w},\lambda}$  has  $D_5$ -contact with  $S$  at  $(0,0,0)$  if  $(0,0)$  is parabolic but not umbilic (i.e.,  $k_i = 0, k_j \neq 0, i \neq j$ ),  $\mathbf{w}$  is parallel to the principal vector  $\mathbf{v}_i$  (i.e.,  $w_j = 0$ ),  $1/\lambda$  is the other principal curvature (i.e.,  $1/\lambda = k_j$ ),  $c_3$  has a double root  $(p_0, q_0)$ , and  $8c_4(p_0, q_0) \neq k_1^3 p_0^4 + k_2^3 q_0^4$ .

**Theorem 4.5.** The cylinder  $C_{\mathbf{w},\lambda}$  has  $A_4$ -contact with  $S$  at  $(0,0,0)$  if and only if one of the following conditions hold.

(A<sub>4a</sub>)  $(0,0)$  is not parabolic (i.e.,  $k_1 k_2 \neq 0$ ),  $\mathbf{w}$  is cylindrical,  $1/\lambda$  is cylindrical curvature with respect to  $\mathbf{w}$ , the critical value set of the orthogonal projection with respect to  $\mathbf{w}$  is nonsingular and the second order vertex (there is a circle with 5-point contact) there.

(A<sub>4b</sub>)  $(0,0)$  is parabolic but not umbilic (i.e.,  $k_i = 0, k_j \neq 0, i \neq j$ ),  $\mathbf{w}$  is parallel to the principal vector  $\mathbf{v}_i$  (i.e.,  $w_j = 0$ ),  $1/\lambda$  is not a principal curvature (i.e.,  $\lambda \neq 1/k_j$ ),  $(0,0)$  is ridge with respect to  $\mathbf{v}_i$ , and

- the line generated by  $\mathbf{w}$  has 4-point contact with  $S$ , the branch  $D'_{\mathbf{w}}$  has  $(2,5)$  cusp and  $1/\lambda$  is the limit of its curvature at  $(0,0)$ , or
- the line generated by  $\mathbf{w}$  has 5-point contact with  $S$ , and  $(0,0)$  is  $\mathbf{v}_j$ -subparabolic.

(A<sub>4c</sub>)  $(0,0)$  is flat umbilic (i.e.,  $k_1 = k_2 = 0$ ),  $c_3(\mathbf{w}) = 0$ ,

$$2c_4(\mathbf{w}) + \lambda \langle J\mathbf{w}, \nabla c_3(\mathbf{w}) \rangle^2 = 0, \text{ and}$$

$$c_5(\mathbf{w}) + \lambda \langle J\mathbf{w}, \nabla c_3(\mathbf{w}) \rangle \langle J\mathbf{w}, \nabla c_4(\mathbf{w}) \rangle + \lambda^2 \langle \mathbf{w}, \nabla c_3(J\mathbf{w}) \rangle \langle J\mathbf{w}, \nabla c_3(\mathbf{w}) \rangle^2 \neq 0.$$

**Theorem 4.6.** The cylinder  $C_{\mathbf{w},\lambda}$  has  $A_5$ -contact with  $S$  at  $(0,0,0)$  if and only if one of the following conditions hold.

(A<sub>5a</sub>)  $(0,0)$  is not parabolic (i.e.,  $k_1 k_2 \neq 0$ ),  $\mathbf{w}$  is cylindrical,  $1/\lambda$  is cylindrical curvature with respect to  $\mathbf{w}$ , the critical value set of the orthogonal projection with respect to  $\mathbf{w}$  is nonsingular and the third order vertex (there is a circle with 6-point contact) there.

(A<sub>5</sub>b)  $k_i = 0, k_j \neq 0, i \neq j, w_j = 0, \lambda \neq 1/k_j, (0, 0)$  is ridge with respect to  $\mathbf{v}_i$ .

- the line generated by  $\mathbf{w}$  has 4-point contact with  $S$ ,  $D'_\mathbf{w}$  has (2, 7) cusp or worse at  $(0, 0)$ ,  $1/\lambda$  is the limit of its curvature at  $(0, 0)$  of  $D'_\mathbf{w}$ , and the derivative of the curvature of  $D'_\mathbf{w}$  by the arc length parameter of  $D'_\mathbf{w}$  does not tend to zero at  $(0, 0)$ , or
- the line generated by  $\mathbf{w}$  has at least 6-point contact with  $S$ , and
  - \*  $a_{60} - 10a_{31}^2(k_2 - 1/\lambda) \neq 0$ , if  $k_1 = 0$ ,
  - \*  $a_{06} - 10a_{13}^2(k_1 - 1/\lambda) \neq 0$ , if  $k_2 = 0$ .

(A<sub>5</sub>c)  $(0, 0)$  is a flat umbilic (i.e.,  $k_1 = k_2 = 0$ ),  $c_3(\mathbf{w}) = 0$ ,

$$\begin{aligned}
& 2c_4(\mathbf{w}) + \lambda \langle J\mathbf{w}, \nabla c_3(\mathbf{w}) \rangle^2 = 0, \\
& c_5(\mathbf{w}) + \lambda \langle J\mathbf{w}, \nabla c_3(\mathbf{w}) \rangle \langle J\mathbf{w}, \nabla c_4(\mathbf{w}) \rangle + \lambda^2 \langle \mathbf{w}, \nabla c_3(J\mathbf{w}) \rangle \langle J\mathbf{w}, \nabla c_3(\mathbf{w}) \rangle^2 = 0, \text{ and} \\
& 2c_6(\mathbf{w}) + \lambda (\langle J\mathbf{w}, \nabla c_4(\mathbf{w}) \rangle^2 / 2 + \langle J\mathbf{w}, \nabla c_5(\mathbf{w}) \rangle \langle J\mathbf{w}, \nabla c_4(\mathbf{w}) \rangle) \\
& + \lambda^2 \langle J\mathbf{w}, \nabla c_3(\mathbf{w}) \rangle (b_{22} \langle J\mathbf{w}, \nabla c_3(\mathbf{w}) \rangle + \langle J\mathbf{w}, \nabla c_4(\mathbf{w}) \rangle \langle \mathbf{w}, \nabla c_3(J\mathbf{w}) \rangle) \\
& + \lambda^3 \langle \mathbf{w}, \nabla c_3(J\mathbf{w}) \rangle^2 (2 \langle \mathbf{w}, \nabla c_3(J\mathbf{w}) \rangle^2 + c_3(J\mathbf{w}) \langle \mathbf{w}, \nabla c_3(J\mathbf{w}) \rangle) \neq 0, \\
& \text{where } b_{22} = \frac{w_2^2}{2} (c_4)_{uu}(\mathbf{w}) - w_1 w_2 (c_4)_{uv}(\mathbf{w}) + \frac{w_1^2}{2} (c_4)_{vv}(\mathbf{w}).
\end{aligned}$$

**Remark 4.7.** Similar to Remark 4.2, there are no  $A_4, A_5$ -contact cylinder at a point in a parabolic line on a generic surface.

## 4.2 Contact map with cylinders and its singularity type

First we show Monge normal form of cylinders  $C_{\mathbf{w}, \lambda}$ .

**Proposition 4.8.** Monge normal form of the cylinder which contains  $(0, 0, 0)$  and whose central axis contains  $(0, 0, \lambda)$  is given by

$$(u, v) \mapsto \left( u, v, \sum_{n=1}^{\infty} (-1)^{n-1} \binom{1/2}{n} \frac{(w_1 v - w_2 u)^{2n}}{\lambda^{2n-1}} \right) \quad (4.1)$$

where  $\mathbf{w}$  denotes a unit vector parallel to the central axis of the cylinder.

*Proof.* Set  $x_1 = w_1 v - w_2 u, y_1 = w_1 u + w_2 v$ . Then  $(x_1, y_1, z)$  form an orthonormal coordinate system, and the cylinder is defined by  $x_1^2 + (z - \lambda)^2 = \lambda^2$ . We thus have

$$z = \lambda \left[ 1 \pm (1 - (x_1/\lambda)^2)^{\frac{1}{2}} \right] = \lambda \left( 1 \pm \left( 1 + \sum_{n=1}^{\infty} (-1)^n \binom{1/2}{n} (x_1/\lambda)^{2n} \right) \right),$$

and we obtain the result. □

We investigate contact of a surface given by (2.1) with the cylinder (4.1). To do this, we consider the difference of the third components of (2.1) and (4.1), namely, the contact map defined by

$$F_{\mathbf{w},\lambda} = f(u, v) - \sum_{n=1}^{\infty} (-1)^{n-1} \binom{1/2}{n} \frac{(w_1 v - w_2 u)^{2n}}{\lambda^{2n-1}}.$$

Remark that Hesse matrix of  $F_{\mathbf{w},\lambda}$  at  $(0, 0)$  is

$$\begin{pmatrix} k_1 - \frac{w_2^2}{\lambda} & \frac{w_1 w_2}{\lambda} \\ \frac{w_1 w_2}{\lambda} & k_2 - \frac{w_1^2}{\lambda} \end{pmatrix}. \quad (4.2)$$

Moreover, the  $k$ -jet of  $F_{\mathbf{w},\lambda}$  at  $(0, 0)$  is

$$\begin{aligned} & \frac{1}{2} \left( (k_1 - \frac{w_2^2}{\lambda}) u^2 + 2 \frac{w_1 w_2}{\lambda} uv + (k_2 - \frac{w_1^2}{\lambda}) v^2 \right) \\ & + \sum_{i+j=3}^k \frac{a_{i,j}}{i!j!} u^i v^j - \sum_{n=2}^{2n \leq k} (-1)^n \binom{1/2}{n} \frac{(-w_2 u + w_1 v)^{2n}}{\lambda^{2n-1}} \end{aligned}$$

**Proposition 4.9.** *The following conditions are equivalent.*

- The function  $F_{\mathbf{w},\lambda}$  has a non-degenerate critical point at  $(0, 0)$ .
- $k_1 k_2 \lambda \neq \kappa_n(\mathbf{w})$ .

*Proof.* They are equivalent that the determinant of (4.2) is not zero.  $\square$

**Corollary 4.10.** *Hesse matrix (4.2) is 0, if and only if*

- $k_1 = 0, k_2 \neq 0, \mathbf{w} = \pm \partial_u, \lambda = 1/k_2$ , or
- $k_2 = 0, k_1 \neq 0, \mathbf{w} = \pm \partial_v, \lambda = 1/k_1$ .

*So Hesse matrix is rank one if one of the following conditions hold.*

- $k_1 k_2 \neq 0, \lambda = \frac{\kappa_n(\mathbf{w})}{k_1 k_2}$ .
- $k_i = 0, k_j \neq 0, w_j = 0, \lambda \neq 1/k_j$  ( $i \neq j$ ).
- $k_1 = k_2 = 0$ .

*When Hesse matrix is of rank 1, its kernel direction is the conjugate direction of  $w_1 \partial_u + w_2 \partial_v$ , which is generated by  $k_2 w_2 \partial_u - k_1 w_1 \partial_v$ , when  $(k_1, k_2) \neq (0, 0)$ . When  $k_1 = k_2 = 0$ , the kernel direction is generated by  $w_2 \partial_u - w_1 \partial_v$ .*

*Proof.* Consequence of the expression (4.2).  $\square$

To show the results in §4.1, it is enough to investigate the conditions that the contact map  $F_{\mathbf{w},\lambda}$  has a singularities of type  $A_2, A_3, A_4, A_5, D_4$  and  $D_5$  for cases (a), (b) and (c). For convenience to refer, we remark that

$$F_{\mathbf{w},\lambda}(u, v) = \begin{cases} \frac{(k_1 w_1 u + k_2 w_2 v)^2}{2 \kappa_n(\mathbf{w})} + o(2) & (k_1 k_2 \neq 0, \lambda = \frac{\kappa_n(\mathbf{w})}{k_1 k_2}), \\ \frac{1}{2} (k_2 - \frac{1}{\lambda}) v^2 + o(2) & (k_1 = 0, \mathbf{w} = (1, 0)), \\ -\frac{1}{2} (w_2 u - w_1 v)^2 + o(2) & (k_1 = k_2 = 0). \end{cases}$$

**4.2.1 Case (a):**  $k_1 k_2 \neq 0$ ,  $\lambda = \frac{\kappa_n(\mathbf{w})}{k_1 k_2}$ .

When  $k_1 k_2 \neq 0$ , we call  $\lambda = \kappa_n(\mathbf{w})/(k_1 k_2)$  **cylindrical radius** and  $1/\lambda$  **cylindrical curvature** with respect to the direction  $\mathbf{w}$ . The cylindrical radius  $\lambda$  is given by

$$\lambda = \frac{w_2^2}{k_1} + \frac{w_1^2}{k_2} = \frac{1 - \cos 2\theta}{2k_1} + \frac{1 + \cos 2\theta}{2k_2} = \frac{1/k_1 + 1/k_2}{2} - \frac{1/k_1 - 1/k_2}{2} \cos 2\theta$$

where  $w_1 = \cos \theta$ ,  $w_2 = \sin \theta$ . We thus have the following properties.

- If  $(0, 0)$  is not umbilic (i.e.,  $k_1 \neq k_2$ ) and is not parabolic point, then the principal directions attain the maximum and the minimum of the cylindrical curvatures.
- If  $(0, 0)$  is umbilic (i.e.,  $k_1 = k_2$ ) and not flat, then the cylindrical curvature does not depend on the tangent vector  $\mathbf{w}$ .

Assume that  $k_1 k_2 \neq 0$ ,  $\kappa_n(\mathbf{w}) \neq 0$  and  $\lambda = \kappa_n(\mathbf{w})/(k_1 k_2)$ . Remark that  $\kappa_n(\mathbf{w}) \neq 0$  since  $\lambda \neq 0$ . We have

$$F_{\mathbf{w}, \lambda} = \frac{(k_1 w_1 u + k_2 w_2 v)^2}{2 \kappa_n(\mathbf{w})} + \sum_{s=3}^k c_s - \sum_{n=2}^{\infty} \frac{(-1)^{n-1}}{\lambda^{2n-1}} (w_1 v - w_2 u)^{2n} + o(k).$$

**Proposition 4.11.**  $F_{\mathbf{w}, \lambda}$  has a singularity of type  $A_2$  at  $(0, 0)$  if  $c_3(\bar{\mathbf{w}}) \neq 0$ .

*Proof.* Setting  $t = k_1 w_1 u + k_2 w_2 v$ , and  $s = w_1 v - w_2 u$ , we have

$$u = \frac{1}{\kappa_n(\mathbf{w})} (w_1 t - k_2 w_2 s), \quad v = \frac{1}{\kappa_n(\mathbf{w})} (w_2 t + k_1 w_1 s).$$

Set  $\hat{c}_s(t, s) = c_s(\frac{1}{\kappa_n(\mathbf{w})} (w_1 t - k_2 w_2 s), \frac{1}{\kappa_n(\mathbf{w})} (w_2 t + k_1 w_1 s))$ . We obtain that

$$F_{\mathbf{w}, \lambda} = \frac{t^2}{2 \kappa_n(\mathbf{w})} + \hat{c}_3(t, s) + o(3).$$

This singularity is of type  $A_2$  if  $\hat{c}_3(0, 1) \neq 0$ . This condition is equivalent that  $c_3(k_2 w_2, -k_1 w_1) \neq 0$  and we conclude the proof.  $\square$

This shows that the contact map  $F_{\mathbf{w}, \lambda}$  has a singularity of type  $A_3$  or worse when  $c_3(\bar{\mathbf{w}}) = 0$ .

Since  $\kappa_n(\mathbf{w})$  is the normal curvature  $k_1 w_1^2 + k_2 w_2^2$  with respect to the direction  $\mathbf{w} = w_1 \partial_u + w_2 \partial_v$ , the cylindrical curvature times the normal curvature  $\kappa_n(\mathbf{w})$  is Gauss curvature  $k_1 k_2$ . This implies the cylindrical curvature is the curvature of the critical value locus of the orthogonal projection to the orthogonal plane to  $\mathbf{w}$  by Koendrink's theorem ([12, Appendix], [13, page 433]).

**Proposition 4.12.** Assume that  $\kappa_n(\mathbf{w}) \neq 0$ . The critical value set has a circle with  $(k + 1)$ -point contact (i.e.,  $(k - 2)$ -th order vertex) if and only if  $F_{\mathbf{w}, \lambda}$  has  $A_k$  singularities at  $(0, 0)$ .

*Proof.* First we remark that the critical locus of the orthogonal projection  $(u, v) \mapsto (w_1v - w_2u, f(u, v))$  is zero of

$$-\det(\mathbf{x}_u \ \mathbf{x}_v \ \mathbf{w}) = w_1f_u + w_2f_v = k_1w_1u + k_2w_2v + \sum_{j=3}^k \langle \mathbf{w}, \nabla c_j \rangle + o(k).$$

Set  $\xi = \langle \mathbf{w}, \nabla f \rangle$ ,  $\eta = w_1v - w_2u$ . Since

$$J = \begin{vmatrix} \frac{\partial \xi}{\partial u} & \frac{\partial \xi}{\partial v} \\ \frac{\partial \eta}{\partial u} & \frac{\partial \eta}{\partial v} \end{vmatrix} = \begin{vmatrix} w_1f_{uu} + w_2f_{uv} & w_1f_{uv} + w_2f_{vv} \\ -w_2 & w_1 \end{vmatrix} = w_1^2f_{uu} + 2w_1w_2f_{uv} + w_2^2f_{vv}$$

$(\xi, \eta)$  form a local coordinate at  $(0, 0)$ , whenever  $J_0 = \kappa_n(\mathbf{w}) \neq 0$ . Since

$$\begin{pmatrix} \frac{\partial u}{\partial \xi} & \frac{\partial u}{\partial \eta} \\ \frac{\partial v}{\partial \xi} & \frac{\partial v}{\partial \eta} \end{pmatrix} = \begin{pmatrix} \frac{\partial \xi}{\partial u} & \frac{\partial \xi}{\partial v} \\ \frac{\partial \eta}{\partial u} & \frac{\partial \eta}{\partial v} \end{pmatrix}^{-1} = \frac{1}{J} \begin{pmatrix} \frac{\partial \eta}{\partial v} & -\frac{\partial \xi}{\partial v} \\ -\frac{\partial \eta}{\partial u} & \frac{\partial \xi}{\partial u} \end{pmatrix} = \frac{1}{J} \begin{pmatrix} w_1 & -w_1f_{uv} - w_2f_{vv} \\ w_2 & w_1f_{uu} + w_2f_{uv} \end{pmatrix},$$

we obtain  $\partial_\xi = \frac{\partial u}{\partial \xi} \partial_u + \frac{\partial v}{\partial \xi} \partial_v = \frac{1}{J}(w_1 \partial_u + w_2 \partial_v)$ . Setting  $f = \frac{1}{2}(k_1u^2 + k_2v^2) + H$ ,  $H_1 = \langle \mathbf{w}, \nabla H \rangle$ ,  $H_2 = w_1^2H_{uu} + 2w_1w_2H_{uv} + w_2^2H_{vv}$ , we obtain  $\partial_\xi H = \frac{H_1}{J}$ ,  $\partial_\xi H_1 = \frac{H_2}{J}$ , and  $J = J_0 + H_2$ .

Since  $\xi = k_1w_1u + k_2w_2v + H_1$ , we have

$$F_{\mathbf{w}, \lambda} = \frac{(\xi - H_1)^2}{2J_0} + H - \sum_{n=2}^{\infty} (-1)^{n-1} \binom{1/2}{n} \eta^{2n},$$

and we obtain that

$$\begin{aligned} \partial_\xi F_{\mathbf{w}, \lambda} &= \frac{(\xi - H_1)(1 - \partial_\xi H_1)}{J_0} + \partial_\xi H = \frac{(\xi - H_1)(1 - H_2/J)}{J_0} + \frac{H_1}{J} \\ &= \frac{(\xi - H_1)(J - H_2)}{J_0 J} + \frac{H_1}{J} = \frac{\xi - H_1}{J} + \frac{H_1}{J} = \frac{\xi}{J}. \end{aligned}$$

Thus  $\partial_\xi F_{\mathbf{w}, \lambda}|_{\xi=0}$  is zero. Now consider Taylor expansion of  $F_{\mathbf{w}, \lambda}$  in the coordinate  $(\xi, \eta)$ :

$$F_{\mathbf{w}, \lambda} = \sum_{i,j} c_{i,j} \xi^i \eta^j.$$

We have  $c_{20} = \frac{1}{2J_0}$  and  $c_{1,j} = 0$  for all  $j$ . So the singularity type of  $F_{\mathbf{w}, \lambda}$  at  $(0, 0)$  is  $A_k$  if and only if  $c_{0,i} = 0$ ,  $i = 1, 2, \dots, k$ ,  $c_{0,k+1} \neq 0$ . This is equivalent that  $F_{\mathbf{w}, \lambda}|_{\xi=0}$  is a unit multiple of  $\eta^{k+1}$ , which means that the critical value set has a circle with  $(k+1)$ -points contact.  $\square$

#### 4.2.2 Case (b): $k_1 = 0$ , $k_2 \neq 0$ , $\mathbf{w} = \partial_u$ .

In this case, we have

$$F_{\mathbf{w}, \lambda} = \frac{1}{2} \left( k_2 - \frac{1}{\lambda} \right) v^2 + h(u, v) + \sum_{n=2}^{\infty} (-1)^n \binom{1/2}{n} \frac{v^{2n}}{\lambda^{2n-1}}$$

where  $h(u, v) = f(u, v) - \frac{1}{2}k_2v^2$ .

Assume that  $\lambda \neq 1/k_2$ . We see the singularity type of  $F_{\mathbf{w}, \lambda}$  is  $A_2$  if  $a_{30} \neq 0$ , since the coefficient of  $u^3$  of  $F_{\mathbf{w}, \lambda}$  is  $a_{30}/6$ . This shows the condition  $(A_2b)$  of Theorem 4.1.

Replacing  $v$  by  $v - a_{21}u^2/2(k_2\lambda - 1)$ , the coefficients of  $u^2v$ ,  $u^4$  become

$$0, \frac{a_{40}}{24} - \frac{a_{21}^2}{8(k_2 - 1/\lambda)},$$

respectively. We see the singularity type of  $F_{\mathbf{w}, \lambda}$  is  $A_3$  if  $a_{40}(k_2 - 1/\lambda) \neq 3a_{21}^2$ . If the line generated by  $\mathbf{w}$  has at least 5-point contact with  $S$  (i.e.,  $a_{40} = 0$ ), then this condition becomes  $a_{21} \neq 0$ , that is,  $(0, 0)$  is not  $\mathbf{v}_2$ -subparabolic. If the line generated by  $\mathbf{w}$  has 4-point contact with  $S$  (i.e.,  $a_{40} \neq 0$ ), then this becomes  $1/\lambda \neq k_2 - 3a_{21}^2/a_{40}$  and the right hand side is the limit of curvature of  $D'_{\mathbf{w}}$  at  $(0, 0)$ . This shows the condition  $(A_3b)$  of Theorem 4.1.

When  $a_{40}(k_2 - 1/\lambda) = 3a_{21}^2$  and  $a_{21} \neq 0$ , the coefficient of  $u^4$  is zero and we see the coefficient of  $u^5$  is

$$\frac{a_{50}}{120} + \frac{a_{40}(a_{40}a_{12} - 2a_{31}a_{21})}{72a_{21}^2}.$$

This shows the condition  $(A_4b)$  of Theorem 4.5.

Replacing  $v$  by  $v - a_{40}(a_{40}a_{12} - a_{31}a_{21})u^3/8a_{21}^3$ , the coefficients of  $u^3v$ ,  $u^6$  become

$$0, \frac{a_{60}a_{21}^4 - 45a_{41}a_{40}a_{21}^3 - 5a_{40}^3a_{21}a_{03} - 30a_{40}^3a_{12}^2 - 30a_{40}a_{31}^2a_{21}^2 + 45a_{40}^2a_{22}a_{21}^2 + 60a_{40}^2a_{31}a_{21}a_{12}}{6480a_{21}^4},$$

respectively. So the singularity type of  $F_{\mathbf{w}, \lambda}$  is  $A_5$  if the later number is non zero.

To complete the proof, we describe the branch  $D'_{\mathbf{w}}$ . A parametrization of the branch of critical set of  $P_{\mathbf{w}}$  whose tangent is generated by  $\mathbf{w}$  is given by  $t \mapsto (\sqrt{\pm 6a_{21}/a_{40}}t + o(1), \pm t^2)$ , and thus  $D'_{\mathbf{w}}$  is parametrized by

$$t \mapsto \left( \mp t^2, \frac{a_{40}k_2 - 3a_{21}^2}{2a_{40}}t^4 + \left[ \mp \frac{3a_{21}}{2a_{40}^5} \right]^{\frac{1}{2}} \frac{3a_{21}^2a_{50} + 5a_{12}a_{40}^2 - 10a_{21}a_{31}a_{40}}{5}t^5 + o(5) \right)$$

whenever  $a_{40} \neq 0$ . This shows the limit of the curvature of  $D'_{\mathbf{w}}$  is  $k_2 - 3a_{21}^2/a_{40}$  at  $(0, 0)$  and  $D'_{\mathbf{w}}$  has  $(2, 5)$  cusp when  $3a_{21}^2a_{50} + 5a_{12}a_{40}^2 - 10a_{21}a_{31}a_{40} \neq 0$ , which shows Condition  $(A_4b)$  in case  $a_{21} \neq 0$ . Remark that the coefficient of  $t^6$  in the second component is

$$\frac{-1}{240a_{21}a_{40}^4} \left[ 8a_{40}(a_{60}a_{21}^4 - 45a_{41}a_{40}a_{41} - 30a_{40}a_{31}^2a_{21}^2 + 45a_{40}^2a_{22}a_{21}^2 + 60a_{40}^2a_{31}a_{21}a_{12}) \right].$$

which shows Condition  $(A_5b)$  in case  $a_{21} \neq 0$ .

When  $a_{40}(k_2 - 1/\lambda) = 3a_{21}^2$  and  $a_{21} = 0$ , we have  $a_{40} = 0$ . Thus we have that

$$F_{\mathbf{w}, \lambda} = \frac{k_2 - 1/\lambda}{2}v^2 + \frac{a_{50}}{120}u^5 + \frac{a_{60} - 10a_{31}^2(k_2 - 1/\lambda)}{720}u^6 + \dots,$$

and the singularity type of  $F_{\mathbf{w}, \lambda}$  is  $A_4$  (resp.  $A_5$ ) if  $a_{50} \neq 0$  (resp.  $a_{50} = 0$ ,  $a_{60} - 10a_{31}^2(k_2 - 1/\lambda) \neq 0$ ).

**Proposition 4.13.** *Assume that  $\lambda = 1/k_2$ .*

- *If  $c_3$  does not have multiple roots, then  $F_{\mathbf{w},\lambda}$  has a singularity of type  $D_4$  at  $(0,0)$ .*
- *If  $c_3$  has a double root  $(p_0, q_0)$  and a single root  $(p_1, q_1)$ , then  $F_{\mathbf{w},\lambda}$  has a singularity of type  $D_5$  at  $(0,0)$ , whenever  $c_4(p_0, q_0) \neq \frac{k_2^3}{8} q_0^4$ .*

*Proof.* When  $\lambda = 1/k_2$ , observe that

$$F_{\mathbf{w},\lambda} = \sum_{s=3}^k c_s(u, v) - \sum_{n=2}^{\infty} (-1)^n \binom{1/2}{n} k_2^{2n-1} v^{2n} + o(k).$$

The first statement is well-known. In the second case we may assume that  $c_3 = (q_0 u - p_1 v)^2 (q_1 u - p_1 v)$ . Setting  $u = -p_1 x + p_0 y$ ,  $v = -q_1 x + q_0 y$ , we have

$$F_{\mathbf{w},\lambda} = \delta^3 x^2 y + [c_4(p_0, q_0) - \frac{k_2^3}{8} q_0^4] y^4 + \dots$$

where  $\delta = p_0 q_1 - q_0 p_1$ . We thus conclude the proof.  $\square$

### 4.2.3 Case (c): $k_1 = k_2 = 0$ .

In this case, we have

$$F_{\mathbf{w},\lambda} = -\frac{(w_1 v - w_2 u)^2}{2\lambda} + f(u, v) - \sum_{n=2}^{\infty} (-1)^{n-1} \binom{1/2}{n} \frac{(w_1 v - w_2 u)^{2n}}{\lambda^{2n-1}}.$$

Setting  $u = w_1 s - w_2 t$ ,  $v = w_2 s + w_1 t$ . we have

$$F_{\mathbf{w},\lambda} = -\frac{t^2}{2\lambda} + f(w_1 s - w_2 t, w_2 s + w_1 t) - \sum_{n=2}^{\infty} (-1)^{n-1} \binom{1/2}{n} \frac{t^{2n}}{\lambda^{2n-1}}.$$

When we write  $f(w_1 s - w_2 t, w_2 s + w_1 t) = \sum_{i,j}^{i+j \leq k} b_{i,j} s^i t^j + o(k)$ , we conclude that

$$b_{30} = c_3(\mathbf{w}), \quad b_{21} = \langle J\mathbf{w}, \nabla c_3(\mathbf{w}) \rangle, \quad b_{12} = \langle \mathbf{w}, \nabla c_3(J\mathbf{w}) \rangle, \quad b_{03} = c_3(J\mathbf{w}), \quad b_{40} = c_4(\mathbf{w}),$$

$$b_{31} = \langle J\mathbf{w}, \nabla c_4(\mathbf{w}) \rangle, \quad b_{22} = \frac{w_2^2}{2} (c_4)_{uu}(\mathbf{w}) - w_1 w_2 (c_4)_{uv}(\mathbf{w}) + \frac{w_1^2}{2} (c_4)_{vv}(\mathbf{w}), \quad \dots,$$

$$b_{50} = c_5(\mathbf{w}), \quad b_{41} = \langle J\mathbf{w}, \nabla c_5(\mathbf{w}) \rangle, \quad \dots, \quad b_{60} = c_6(\mathbf{w}), \quad \dots, \quad \text{where } J\mathbf{w} = w_2 \partial_u - w_1 \partial_v.$$

The condition  $(A_{2c})$  of Theorem 4.1 follows to see the coefficient  $b_{30}$ .

Replacing  $t$  by  $t + \lambda b_{21} s^2$ , the coefficient of  $s^2 t$  and  $s^4$  become 0,  $b_{40} + \lambda b_{21}^2 / 2$ , respectively. The condition  $(A_{3c})$  follows from Theorem 4.1.

Observe the coefficient of  $s^5$  is  $b_{50} + \lambda b_{31} b_{21} + \lambda^2 b_{12} b_{21}^2$ , which shows the condition  $(A_{4c})$  of Theorem 4.5.

Replacing  $t$  by  $t + \lambda(b_{31} + b_{21}^2 \lambda) s^3$ , we obtain that the coefficients of  $s^3 t$  and  $s^6$  become

$$0, \quad b_{60} + \lambda(b_{31}^2 / 2 + b_{41} b_{21}) + \lambda^2 b_{21} (b_{22} b_{21} + 2b_{31} b_{12}) + \lambda^3 b_{21}^2 (2b_{12}^2 + b_{03} b_{21}),$$

respectively. This concludes the condition  $(A_{5c})$  of Theorem 4.6

## 5 Cylindrical directions

In Theorem 4.1, we observe that the kernel field of  $A_{\geq 3}$ -contact with cylinders is cylindrical directions whenever  $k_1 k_2 \neq 0$ . These are also defined at nearby points, and we express it in terms of Monge normal form. This field is defined by  $\bar{C}(du, dv) = 0$  where  $\bar{C}(du, dv) = C(\kappa_2 dv, -\kappa_1 du)$ , and this can be considered as a counterpart of principal fields. Here  $C(p, q)$  denote Monge cubic defined in Proposition 5.2. This section is dedicated to analysis of this field.

Let  $[p : q]$  denote the homogeneous coordinate of the real projective line  $P^1$ . Let  $M$  denote the subset defined by  $C(p, q) = 0$  in  $\mathbb{R}^2 \times P^1$ ,  $\pi : M \rightarrow \mathbb{R}^2$  the natural projection, and  $\bar{\omega} = p\kappa_1 du + q\kappa_2 dv$ .

**Theorem 5.1.** *Assume that  $(0, 0)$  is not an umbilic of  $\mathbf{x}$  (i.e.,  $k_1 \neq k_2$ ). The flows of  $\bar{C}(du, dv) = 0$  is obtained by the projection image of the flows of 1-form  $\bar{\omega}|_M$  by the natural projection  $M \rightarrow \mathbb{R}^2$ . The 1-form  $\bar{\omega}|_M$  is singular at  $(0, 0) \times [p_0, q_0] \in M$  if and only if one of the following conditions holds.*

1.  $(p_0, q_0)$  is a single root of  $C(p, q) = 0$  at  $(u, v) = (0, 0)$ , and
  - (a)  $\kappa_1(0, 0) = 0$ ,  $(0, 0)$  is ridge with respect to  $\mathbf{v}_1$ , and  $(p_0, q_0)$  represents the principal direction  $\mathbf{v}_1$  (i.e.,  $a_{3,0} = 0$ ,  $q_0 = 0$ ), or
  - (b)  $\kappa_2(0, 0) = 0$ ,  $(0, 0)$  is ridge with respect to  $\mathbf{v}_2$ , and  $(p_0, q_0)$  represents the principal direction  $\mathbf{v}_2$  (i.e.,  $a_{0,3} = 0$ ,  $p_0 = 0$ ).
2.  $(p_0, q_0)$  is a multiple root of  $C(p, q) = 0$  at  $(u, v) = (0, 0)$ , that is, is in the discriminant set  $D$  of  $C(p, q) = 0$  (i.e.,  $C_p(p_0, q_0) = C_q(p_0, q_0) = 0$ ), and
  - (a)  $D$  is nonsingular at  $(0, 0)$  and the tangent direction of  $D$  at  $(0, 0)$  is conjugate to  $(p_0, q_0)$ , or
  - (b)  $\alpha = \beta = 0$  (in particular,  $D$  is singular at  $(0, 0)$ ) where

$$\alpha = C_u(p_0, q_0)|_{(u,v)=(0,0)}, \quad \beta = C_v(p_0, q_0)|_{(u,v)=(0,0)}.$$

We conclude that the position of singular points of  $\bar{\omega}|_M$  is determined by 3-jet (resp. 4-jet) of Monge normal form (2.1) in Case 1 (resp. Case 2). In generic context, these singularity types are saddle, node or focus, which are determined by 4-jet (resp. 5-jet) of Monge normal form (2.1) in Case 1 (resp. Case 2). The phase portraits of singularities of the flows of the equation  $\bar{C}(du, dv) = 0$ , in the generic context, will be also given. See pictures after Propositions 5.7, 5.11, and Remark 5.9.

### 5.1 Monge normal form of a surface given by Monge normal form

We consider the Monge normal form of the surface defined by (2.1) at the point  $(u, v, f(u, v))$ .

**Proposition 5.2.** *Monge normal form of  $\mathbf{x}$  at  $(u, v, f(u, v))$  is given by*

$$z = \kappa_1 \frac{x^2}{2} + \kappa_2 \frac{y^2}{2} + C(x, y) + o(x, y)^3 \quad (5.1)$$

where

$$\begin{aligned} \kappa_1 &= k_1 + c_{3,uu}(u, v) + c_{4,uu}(u, v) - (3k_1^2 u^2 + k_2^2 v^2) + \frac{2c_{3,uv}(u, v)^2}{k_1 - k_2} + o(u, v)^2, \\ \kappa_2 &= k_2 + c_{3,vv}(u, v) + c_{4,vv}(v, v) - (k_1^2 u^2 + 3k_2^2 v^2) - \frac{2c_{3,uv}(u, v)^2}{k_1 - k_2} + o(u, v)^2, \\ C(x, y) &= c_3(x, y) + c_{4,u}(x, y)u + c_{4,v}(x, y)v + \frac{c_{3,uv}(u, v)}{2(k_1 - k_2)}C_1 \\ &\quad + c_{5,uuu}(u, v)\frac{x^3}{6} + c_{5,uuv}(u, v)\frac{x^2y}{2} + c_{5,uvv}(u, v)\frac{xy^2}{2} + c_{5,vvv}(u, v)\frac{y^3}{6} \\ &\quad + \frac{c_{3,uv}(u, v)}{4(k_1 - k_2)^2}(Pu + Qv) - \frac{Y_1}{4} + \frac{X + Y_2}{4(k_1 - k_2)} - (k_1^2 ux + k_2^2 vy)\frac{k_1 x^2 + k_2 y^2}{2} \\ &\quad + o(u, v)^2, \\ C_1 &= a_{21}x^3 + (2a_{12} - a_{30})x^2y + (a_{03} - 2a_{21})xy^2 - a_{12}y^3 \\ P &= a_{21}(4a_{12} - 3a_{30})x^3 + (4a_{12}^2 + 2a_{03}a_{21} - 7a_{21}^2 - 6a_{12}a_{30} + 2a_{30}^2)x^2y \\ &\quad + (2a_{03}a_{12} - 11a_{12}a_{21} - 2a_{03}a_{30} + 6a_{21}a_{30}xy^2 + (-2a_{12}^2 - a_{03}a_{21} + 2a_{21}^2 + 2a_{12}a_{30})y^3, \\ Q &= (2a_{12}^2 + 2a_{03}a_{21} - 2a_{21}^2 - a_{12}a_{30})x^3 + (6a_{03}a_{12} - 11a_{12}a_{21} - 2a_{03}a_{30} + 2a_{21}a_{30})x^2y \\ &\quad + (2a_{03}^2 - 7a_{12}^2 - 6a_{03}a_{21} + 4a_{21}^2 + 2a_{12}a_{30})xy^2 - a_{12}(3a_{03} - 4a_{21})y^3, \\ X &= [3a_{21}a_{31}u^2 + 2(2a_{21}a_{22} + a_{12}a_{31})uv + (a_{13}a_{21} + 2a_{12}a_{22})v^2]x^3 \\ &\quad + \left[ \begin{array}{l} (4a_{21}a_{22} + 2a_{12}a_{31} - a_{30}a_{31} - 2a_{21}a_{40})u^2 \\ + 2(2a_{13}a_{21} + 4a_{12}a_{22} - a_{22}a_{30} - a_{21}a_{31}) \\ - a_{12}a_{40} \end{array} \right] x^2y + \left[ \begin{array}{l} (2a_{13}a_{21} + a_{03}a_{31} - 6a_{21}a_{31})u^2 + 2(a_{12}a_{13}) \\ + a_{04}a_{21} + a_{03}a_{22} - 4a_{21}a_{22} - 2a_{12}a_{31} \end{array} \right] uv \\ &\quad + [(2a_{04}a_{12} + a_{03}a_{13} - 2a_{13}a_{21} - 4a_{12}a_{22})v^2] xy^2 \\ &\quad + [(-2a_{21}a_{22} - a_{12}a_{31})u^2 - 2(a_{13}a_{21} + 2a_{12}a_{22})uv - 3a_{12}a_{13}v^2]y^3, \\ Y_1 &= 19a_{30}k_1^2 u^2 x^3 + 3(3a_{03}k_1 k_2 + 4a_{21}k_2^2)v^2 x^2 y + 3(4a_{12}k_1^2 + 3a_{30}k_1 k_2)u^2 xy^2 + 19a_{03}k_2^2 v^2 y^3, \\ Y_2 &= [-6a_{21}k_1^2(3k_1 - k_2)uv + (-3a_{12}k_1^3 - 3a_{12}k_1^2 k_2 - a_{30}k_1 k_2^2 + a_{30}k_2^3)v^2]x^3 \\ &\quad + [-3a_{21}k_1(3k_1^2 - 4k_1 k_2 - k_2^2)u^2 + 6k_2(3a_{12}k_1^2 - 2a_{12}k_1 k_2 - a_{30}k_2^2)uv]x^2 y \\ &\quad + [-6k_1(a_{03}k_1^2 + 2a_{21}k_1 k_2 - 3a_{21}k_2^2)uv - 3a_{12}k_2(k_1^2 + 4k_1 k_2 - 3k_2^2)v^2]xy^2 \\ &\quad + [(-a_{03}k_1^3 + a_{03}k_1^2 k_2 + 3a_{21}k_1 k_2^2 + 3a_{21}k_2^3)u^2 - 6a_{12}(k_1 - 3k_2)k_2^2 uv]y^3. \end{aligned}$$

Remark that

- 1(resp. 2)-jet of  $\kappa_i$  is determined by 3(resp. 4)-jet of Monge normal form (2.1), and
- 1(resp. 2)-jet of  $C$  is determined by 4(resp. 5)-jet of Monge normal form (2.1).

We call the cubic form  $C(x, y)$  **Monge cubic**. To show this proposition, we define an orthonormal frame by

$$\mathbf{e}_1(u, v) = \frac{\mathbf{x}_u(u, v)}{\|\mathbf{x}_u(u, v)\|},$$

$$\begin{aligned}\mathbf{e}_2(u, v) &= \frac{[\mathbf{x}_u(u, v) \times \mathbf{x}_u(u, v)] \times \mathbf{x}_u(u, v)}{\|[\mathbf{x}_u(u, v) \times \mathbf{x}_v(u, v)] \times \mathbf{x}_u(u, v)\|}, \\ \mathbf{e}_3(u, v) &= \frac{\mathbf{x}_u(u, v) \times \mathbf{x}_v(u, v)}{\|\mathbf{x}_u(u, v) \times \mathbf{x}_v(u, v)\|}.\end{aligned}$$

These can be written in the following explicit forms:

$$\begin{aligned}\mathbf{e}_1(u, v) &= \frac{1}{\sqrt{1 + f_u(u, v)^2}}(1, 0, f_u(u, v)), \\ \mathbf{e}_2(u, v) &= \frac{(-f_u(u, v)f_v(u, v), 1 + f_u(u, v)^2, f_v(u, v))}{\sqrt{1 + f_u(u, v)^2}\sqrt{1 + f_u(u, v)^2 + f_v(u, v)^2}} \\ \mathbf{e}_3(u, v) &= \frac{(-f_u(u, v), -f_v(u, v), 1)}{\sqrt{1 + f_u(u, v)^2 + f_v(u, v)^2}}\end{aligned}$$

Setting

$$\begin{aligned}\bar{x} &= \langle \mathbf{x}(u + \bar{u}, v + \bar{v}) - \mathbf{x}(u, v), \mathbf{e}_1(u, v) \rangle, \\ \bar{y} &= \langle \mathbf{x}(u + \bar{u}, v + \bar{v}) - \mathbf{x}(u, v), \mathbf{e}_2(u, v) \rangle, \\ z &= \langle \mathbf{x}(u + \bar{u}, v + \bar{v}) - \mathbf{x}(u, v), \mathbf{e}_3(u, v) \rangle,\end{aligned}$$

we obtain

$$\bar{x} = \frac{\bar{u} + [f(u + \bar{u}, v + \bar{v}) - f(u, v)]f_u(u, v)}{\sqrt{1 + f_u(u, v)^2}}, \quad (5.2)$$

$$\bar{y} = \frac{-\bar{u}f_u(u, v)f_v(u, v) + \bar{v}[1 + f_u(u, v)^2] + [f(u + \bar{u}, v + \bar{v}) - f(u, v)]f_v(u, v)}{\sqrt{1 + f_u(u, v)^2}\sqrt{1 + f_u(u, v)^2 + f_v(u, v)^2}}, \quad (5.3)$$

$$z = \frac{-\bar{u}f_u(u, v) - \bar{v}f_v(u, v) + [f(u + \bar{u}, v + \bar{v}) - f(u, v)]}{\sqrt{1 + f_u(u, v)^2 + f_v(u, v)^2}}. \quad (5.4)$$

By inverse mapping theorem,  $(\bar{u}, \bar{v})$  are determined by  $(\bar{x}, \bar{y})$  when  $|(\bar{u}, \bar{v})| < \varepsilon \ll 1$ .

**Lemma 5.3.** *Setting  $f_{i,j} = f_{u^i v^j}(u, v)$ , we have*

$$\begin{aligned}\bar{x} &= \sqrt{1 + f_{10}^2} \bar{u} + \frac{f_{10}f_{01}}{\sqrt{1 + f_{10}^2}} \bar{v} + \frac{f_{10}}{\sqrt{1 + f_{10}^2}} \sum_{i+j \geq 2}^k \frac{f_{i,j}}{i!j!} \bar{u}^i \bar{v}^j + o(\bar{u}, \bar{v})^k, \\ \bar{y} &= \frac{\sqrt{1 + f_{10}^2 + f_{01}^2}}{\sqrt{1 + f_{10}^2}} \bar{v} + \frac{f_{01}}{\sqrt{1 + f_{10}^2}\sqrt{1 + f_{10}^2 + f_{01}^2}} \sum_{i+j \geq 2}^k \frac{f_{i,j}}{i!j!} \bar{u}^i \bar{v}^j + o(\bar{u}, \bar{v})^k.\end{aligned}$$

*Proof.* By (5.2) and (5.3), we have

$$\bar{x}_{\bar{u}} = \frac{1 + f_u(u + \bar{u}, v + \bar{v})f_u(u, v)}{\sqrt{1 + f_u(u, v)^2}}, \quad \bar{y}_{\bar{u}} = \frac{f_v(u, v)(-f_u(u, v) + f_u(u + \bar{u}, v + \bar{v}))}{\sqrt{1 + f_u(u, v)^2}\sqrt{1 + f_u(u, v)^2 + f_v(u, v)^2}},$$

$$\bar{x}_{\bar{v}} = \frac{f_v(u+\bar{u},v+\bar{v})f_u(u,v)}{\sqrt{1+f_u(u,v)^2}}, \quad \bar{y}_{\bar{v}} = \frac{[1+f_u(u,v)^2]+f_v(u+\bar{u},v+\bar{v})f_v(u,v)}{\sqrt{1+f_u(u,v)^2}\sqrt{1+f_u(u,v)^2+f_v(u,v)^2}},$$

and, for  $(i, j)$  with  $i + j \geq 2$ ,

$$\bar{x}_{\bar{u}^i \bar{v}^j} = \frac{f_{u^i v^j}(u+\bar{u},v+\bar{v})f_u(u,v)}{\sqrt{1+f_u(u,v)^2}}, \quad \bar{y}_{\bar{u}^i \bar{v}^j} = \frac{f_{u^i v^j}(u+\bar{u},v+\bar{v})f_v(u,v)}{\sqrt{1+f_u(u,v)^2}\sqrt{1+f_u(u,v)^2+f_v(u,v)^2}}.$$

These imply the lemma.  $\square$

**Lemma 5.4.** *We have*

$$\begin{aligned} \bar{u} &= \frac{1}{\sqrt{1+f_{10}^2}}\bar{x} - \frac{f_{10}f_{01}}{\sqrt{1+f_{10}^2}\sqrt{1+f_{10}^2+f_{01}^2}}\bar{y} \\ &\quad - \frac{f_{10}}{(1+f_{10}^2)(1+f_{10}^2+f_{01}^2)}\left(\frac{f_{11}}{2}\bar{x}^2 + \frac{(1+f_{10}^2)f_{11}-f_{10}f_{01}f_{20}}{\sqrt{1+f_{10}^2+f_{01}^2}}\bar{x}\bar{y}\right. \\ &\quad \left.+ \frac{f_{10}^2f_{01}^2f_{20}-2f_{10}f_{01}(1+f_{10}^2)f_{11}+(1+f_{10}^2)f_{02}}{2(1+f_{10}^2+f_{01}^2)}\bar{y}^2\right) + o(\bar{x}, \bar{y})^2, \\ \bar{v} &= \frac{\sqrt{1+f_{10}^2}}{\sqrt{1+f_{10}^2+f_{01}^2}}\bar{y} \\ &\quad + \frac{f_{01}}{(1+f_{10}^2)(1+f_{10}^2+f_{01}^2)}\left(\frac{f_{20}}{2}\bar{x}^2 + \frac{-f_{10}f_{01}f_{20}+(1+f_{10}^2)f_{11}}{\sqrt{1+f_{10}^2+f_{01}^2}}\bar{x}\bar{y}\right. \\ &\quad \left.+ \frac{f_{10}^2f_{01}^2f_{20}-2(1+f_{10}^2)f_{11}+(1+f_{10}^2)^2f_{02}}{2(1+f_{10}^2+f_{01}^2)}\bar{y}^2\right) + o(\bar{x}, \bar{y})^2. \end{aligned}$$

*Proof.* When we write

$$\bar{x} = \sum_{i+j \geq 1}^k x_{i,j} \frac{\bar{u}^i \bar{v}^j}{i!j!} + o(\bar{u}, \bar{v})^k, \quad \bar{y} = \sum_{i+j \geq 1}^k y_{i,j} \frac{\bar{u}^i \bar{v}^j}{i!j!} + o(\bar{u}, \bar{v})^k, \quad (5.5)$$

$$\bar{u} = \sum_{i+j \geq 1}^k \bar{u}_{i,j} \frac{\bar{x}^i \bar{y}^j}{i!j!} + o(\bar{u}, \bar{v})^k, \quad \bar{v} = \sum_{i+j \geq 1}^k \bar{v}_{i,j} \frac{\bar{x}^i \bar{y}^j}{i!j!} + o(\bar{u}, \bar{v})^k, \quad (5.6)$$

we obtain that

$$\begin{aligned} u_{10} &= \frac{y_{01}}{x_{10}y_{01} - x_{01}y_{10}} & v_{10} &= \frac{-y_{10}}{x_{10}y_{01} - x_{01}y_{10}} \\ u_{01} &= \frac{-x_{01}}{x_{10}y_{01} - x_{01}y_{10}} & v_{01} &= \frac{x_{10}}{x_{10}y_{01} - x_{01}y_{10}}. \end{aligned}$$

Moreover, we have

$$\begin{pmatrix} x_{10}I_3 & x_{01}I_3 \\ y_{10}I_3 & y_{01}I_3 \end{pmatrix} \begin{pmatrix} U_2 \\ V_2 \end{pmatrix} + \begin{pmatrix} \Delta_2 & 0 \\ 0 & \Delta_2 \end{pmatrix} \begin{pmatrix} X_2 \\ Y_2 \end{pmatrix} = 0$$

where

$$X_2 = \begin{pmatrix} x_{20} \\ x_{11} \\ x_{02} \end{pmatrix}, \quad Y_2 = \begin{pmatrix} y_{20} \\ y_{11} \\ y_{02} \end{pmatrix}, \quad U_2 = \begin{pmatrix} u_{20} \\ u_{11} \\ u_{02} \end{pmatrix}, \quad V_2 = \begin{pmatrix} v_{20} \\ v_{11} \\ v_{02} \end{pmatrix},$$

$$\Delta_2 = \begin{pmatrix} u_{10}^2 & 2u_{10}v_{10} & v_{10}^2 \\ u_{10}u_{01} & u_{10}v_{01} + u_{01}v_{10} & v_{10}v_{01} \\ u_{01}^2 & 2u_{01}v_{01} & v_{01}^2 \end{pmatrix}.$$

Since  $v_{10} = y_{10} = 0$ , we obtain that

$$u_{20} = \frac{x_{01}y_{20} - x_{20}y_{01}}{x_{10}^3 y_{01}}, \quad u_{11} = \frac{-x_{10}x_{11}y_{01} + x_{01}x_{20}y_{01} + x_{01}x_{10}y_{11} - x_{01}^2 y_{20}}{x_{10}^3 y_{01}^2},$$

$$u_{02} = \frac{-x_{02}x_{10}^2 y_{01} + x_{01}(2x_{10}x_{11}y_{01} - x_{01}x_{20}y_{01} + x_{10}^2 y_{02} - 2x_{01}x_{10}y_{11} + x_{01}^2 y_{20})}{x_{10}^3 y_{01}^3},$$

$$v_{20} = \frac{-y_{20}}{x_{10}^2 y_{01}}, \quad v_{11} = \frac{x_{01}y_{20} - x_{10}y_{11}}{x_{10}^2 y_{01}^2}, \quad v_{02} = -\frac{x_{10}^2 y_{02} - 2x_{01}x_{10}y_{11} + x_{01}^2 y_{20}}{x_{10}^2 y_{01}^3}.$$

These conclude the proof.  $\square$

*Proof of Proposition 5.2.* By Taylor's theorem, we have

$$f(u + \bar{u}, v + \bar{v}) - f(u, v) = \sum_{i+j \geq 1}^k \frac{f_{i,j}}{i!j!} \bar{u}^i \bar{v}^j + o(\bar{u}, \bar{v})^k,$$

and thus, by (5.4),

$$z = \frac{1}{\sqrt{1 + f_{10}^2 + f_{01}^2}} \sum_{i+j \geq 2}^k \frac{f_{i,j}}{i!j!} \bar{u}^i \bar{v}^j + o(\bar{u}, \bar{v})^k.$$

By (5.6) and  $v_{10} = 0$ , we have

$$z = \frac{1}{\sqrt{1 + f_{10}^2 + f_{01}^2}} \left( \frac{f_{20}u_{10}^2}{2} \bar{x}^2 + u_{10}(u_{01}f_{20} + v_{01}f_{11}) \bar{x}\bar{y} + \frac{u_{01}^2 f_{20} + 2u_{01}v_{01}f_{11} + v_{01}^2 f_{02}}{2} \bar{y}^2 \right.$$

$$+ \frac{u_{10}(f_{3,0}u_{10}^2 + 3f_{20}u_{20} + 3f_{11}v_{20})}{6} \bar{x}^3$$

$$+ \frac{f_{3,0}u_{01}u_{10}^2 + f_{2,1}u_{10}^2 v_{01} + f_{20}(2u_{10}u_{11} + u_{01}u_{20}) + f_{11}(u_{20}v_{01} + 2u_{10}v_{11} + u_{01}v_{20}) + 2f_{02}v_{01}v_{20}}{2} \bar{x}^2 \bar{y}$$

$$+ \frac{f_{3,0}u_{01}^2 u_{10} + 2f_{2,1}u_{01}u_{10}v_{01} + f_{1,2}u_{10}v_{01}^2}{f_{20}(u_{02}u_{10} + 2u_{01}u_{11}) + f_{11}(2u_{11}v_{01} + u_{10}v_{02} + 2u_{01}v_{11}) + 4f_{02}v_{01}v_{11}} \bar{x}\bar{y}^2$$

$$+ \left. \frac{f_{3,0}u_{01}^3 + 3f_{2,1}u_{01}^2 v_{01} + 3f_{1,2}u_{01}v_{01}^2 + f_{0,3}v_{01}^3 + 3f_{20}u_{01}u_{02} + 3f_{11}(u_{02}v_{01} + u_{01}v_{02}) + 6f_{02}v_{01}v_{02}}{6} \bar{y}^3 \right)$$

$$+ o(\bar{x}, \bar{y})^3$$

We take an orthonormal matrix

$$T = \left( 1 - \frac{c_{3,uv}(u, v)^2}{2(k_1 - k_2)^2} + o(u, v)^2 \right) \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + c \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$$

where  $c = \frac{c_{3,uv}(u,v)}{k_1-k_2} + \frac{a_{2,1}a_{1,2}(u^2-v^2)+(a_{2,1}^2-a_{1,2}^2+a_{3,0}a_{1,2}-a_{2,1}a_{0,3})uv}{(k_1-k_2)^2} + \frac{c_{4,uv}(u,v)-k_1^2k_2}{(k_1-k_2)} + o(u,v)^2$ .  
Composing the rotation

$$\begin{pmatrix} \bar{x} \\ \bar{y} \end{pmatrix} = T \begin{pmatrix} x \\ y \end{pmatrix},$$

we obtain the result.  $\square$

## 5.2 Monge cubic

We assume that  $k_1 \neq k_2$  and that Monge cubic is expressed in a neighbourhood of  $(0,0)$  as follows:

$$C(x,y) = A_{3,0}x^3 + 3A_{2,1}x^2y + 3A_{1,2}xy^2 + A_{0,3}y^3. \quad (5.7)$$

Remember that  $D$  denote the discriminant set of  $C$ , that is, the zero set of

$$\Delta = \begin{vmatrix} A_{3,0} & 2A_{2,1} & A_{1,2} & 0 \\ A_{2,1} & 2A_{1,2} & A_{0,3} & 0 \\ 0 & A_{3,0} & 2A_{2,1} & A_{1,2} \\ 0 & A_{2,1} & 2A_{1,2} & A_{0,3} \end{vmatrix}. \quad (5.8)$$

Let  $\Sigma(M)$  denote the singular set of  $M$ .

**Lemma 5.5.** *The following conditions are equivalent.*

- $(0,0) \times [p_0 : q_0] \in \Sigma(M)$ .
- $C_u = C_v = C_p = C_q = 0$  at  $(0,0) \times [p_0 : q_0]$ , that is,  $\alpha = \beta = 0$ ,

$$A_{3,0}p_0^2 + 2A_{2,1}p_0q_0 + A_{1,2}q_0^2 = A_{2,1}p_0^2 + 2A_{1,2}p_0q_0 + A_{0,3}q_0^2 = 0 \quad (5.9)$$

at  $(0,0)$ .

*Proof.* A consequence of the implicit function theorem.  $\square$

The second condition in the lemma holds for some  $[p_0 : q_0]$  if and only if

$$\text{rank} \begin{pmatrix} \partial_u A_{3,0} & 3\partial_u A_{2,1} & 3\partial_u A_{1,2} & \partial_u A_{0,3} & 0 \\ \partial_v A_{3,0} & 3\partial_v A_{2,1} & 3\partial_v A_{1,2} & \partial_v A_{0,3} & 0 \\ 0 & \partial_u A_{3,0} & 3\partial_u A_{2,1} & 3\partial_u A_{1,2} & \partial_u A_{0,3} \\ 0 & \partial_v A_{3,0} & 3\partial_v A_{2,1} & 3\partial_v A_{1,2} & \partial_v A_{0,3} \\ A_{3,0} & 2A_{2,1} & A_{1,2} & 0 & 0 \\ A_{2,1} & 2A_{1,2} & A_{0,3} & 0 & 0 \\ 0 & A_{3,0} & 2A_{2,1} & A_{1,2} & 0 \\ 0 & A_{2,1} & 2A_{1,2} & A_{0,3} & 0 \\ 0 & 0 & A_{3,0} & 2A_{2,1} & A_{1,2} \\ 0 & 0 & A_{2,1} & 2A_{1,2} & A_{0,3} \end{pmatrix} < 5 \quad \text{at } (0,0), \quad (5.10)$$

because of the result of [11]. Let  $I$  denote the ideal generated by maximal minors of

$$\begin{pmatrix} y_0 & y_1 & y_2 & y_3 & 0 \\ z_0 & z_1 & z_2 & z_3 & 0 \\ 0 & y_0 & y_1 & y_2 & y_3 \\ 0 & z_0 & z_1 & z_2 & z_3 \\ x_0 & 2x_1 & x_2 & 0 & 0 \\ x_1 & 2x_2 & x_3 & 0 & 0 \\ 0 & x_0 & 2x_1 & x_2 & 0 \\ 0 & x_1 & 2x_2 & x_3 & 0 \\ 0 & 0 & x_0 & 2x_1 & x_2 \\ 0 & 0 & x_1 & 2x_2 & x_3 \end{pmatrix}$$

in the ring  $R = \mathbb{R}[x_i, y_i, z_i; i = 0, 1, 2, 3]$ . By computation using Singular, we see the height of  $I$  is 3. This means that, for a generic choice of the coefficients in Monge normal form, we do not have (5.10).

We assume that the rank of the matrix in (5.10) is 5. To investigate the equation  $C(du, dv) = 0$ , we consider the foliation defined by the restriction of the 1-form  $\omega = q du - p dv$  to  $M$ . The singularity of  $\omega|_M$  is defined by  $pC_u + qC_v = 0$ ,  $C_p = C_q = 0$ . Thus the induced flow on  $M$  is singular at  $(0, 0) \times [p_0 : q_0]$  if and only if (5.9) holds, and

$$\alpha p_0 + \beta q_0 = 0 \quad \text{at } (0, 0). \quad (5.11)$$

Remember that  $D$  is the discriminant set of the cubic  $C$  which is zero of  $\Delta$  defined by (5.8). We assume  $(0, 0) \in D$ . Then there is a non-zero  $(p_0, q_0)$  with (5.9), i.e.,

$$p_0^2(p_0 A_{3,0} + q_0 A_{2,1}) = -p_0 q_0(p_0 A_{2,1} + q_0 A_{1,2}) = q_0^2(p_0 A_{1,2} + q_0 A_{0,3}) \quad (5.12)$$

at  $(0, 0)$ .

Let  $\Sigma(D)$  denote the singular set of  $D$  and let  $\Sigma_1$  denote the set defined by

$$\text{rank} \begin{pmatrix} A_{3,0} & A_{2,1} & A_{1,2} \\ A_{2,1} & A_{1,2} & A_{0,3} \end{pmatrix} < 2. \quad (5.13)$$

Remark that (5.13) holds if and only if (5.12) is zero for some non-zero  $(p_0, q_0)$ .

**Lemma 5.6.** • *If  $(0, 0) \in D \setminus \Sigma(D)$ , we have  $(\alpha, \beta) \neq (0, 0)$ , and  $\beta \partial_u - \alpha \partial_v$  is tangent to  $D$  at  $(0, 0)$ .*

- $\Sigma(D) = \Sigma_1 \cup \pi(\Sigma(M))$ .
- *Suppose that  $(0, 0) \in \Sigma(D) \setminus \pi(\Sigma(M))$ . Then, the singularity type of  $\Sigma(D)$  at  $(0, 0)$  is cusp if and only if*

$$\begin{vmatrix} \partial_u A_{3,0} & \partial_u A_{2,1} & \partial_u A_{1,2} & \partial_u A_{0,3} \\ \partial_v A_{3,0} & \partial_v A_{2,1} & \partial_v A_{1,2} & \partial_v A_{0,3} \\ q_0^2 & -2p_0 q_0 & p_0^2 & 0 \\ 0 & q_0^2 & -2p_0 q_0 & p_0^2 \end{vmatrix} + 2p_0^2 q_0^2 A \neq 0 \quad \text{at } (0, 0) \quad (5.14)$$

where  $A = -\partial_u A_{3,0} \partial_v A_{0,3} + \partial_u A_{2,1} \partial_v A_{1,2} - \partial_v A_{2,1} \partial_u A_{1,2} + \partial_u A_{0,3} \partial_v A_{3,0}$ .

*Proof.* Since the linear terms of (5.8) is

$$-\frac{4(p_0q_1 - p_1q_0)^3}{27}(\alpha u + \beta v) \quad (5.15)$$

we have the first assertion.  $(0, 0) \in \Sigma(D)$  if and only if (5.13) or  $(\alpha, \beta) = (0, 0)$  hold. When  $(\alpha, \beta) = (0, 0)$  holds, then we have  $(0, 0) \times [p_0 : q_0] \in \Sigma(M)$  by (5.9). This shows the second result. We assume that (5.13) holds. Then the constant term of  $C$  is a constant multiple of  $(q_0du - p_0dv)^3$ . We observe that the quadric part of (5.8) is

$$-(\alpha u + \beta v)^2$$

and the kernel direction for Hesse matrix is generated by  $\beta\partial_u - \alpha\partial_v$ . If we evaluate the cubic part of (5.8) over this vector, we obtain a constant multiple of the left hand side of the cube of (5.14), which shows the last result.  $\square$

### 5.3 Conjugate Monge cubic

Assume that  $k_1 \neq k_2$ . Let  $\bar{C}(p, q)$  denote the **conjugate Monge cubic** defined by  $\bar{C}(p, q) = C(\kappa_2q, -\kappa_1p)$ , i.e.,

$$\bar{C}(p, q) = A_{3,0}\kappa_2^3q^3 - 3A_{2,1}\kappa_1\kappa_2^2pq^2 + 3A_{1,2}\kappa_1^2\kappa_2p^2q - A_{0,2}\kappa_1^3p^3.$$

We call  $\bar{C}(du, dv) = 0$  the **cylindrical cubic differential equation**. Let  $\bar{D}$  denote the discriminant of  $\bar{C}$ , that is, the zero of

$$\bar{\Delta} = \begin{vmatrix} -A_{0,3}\kappa_1^3 & 2A_{1,2}\kappa_1^2\kappa_2 & -A_{2,1}\kappa_1\kappa_2^2 & 0 \\ A_{1,2}\kappa_1^2\kappa_2 & -2A_{2,1}\kappa_1\kappa_2^2 & A_{3,0}\kappa_2^3 & 0 \\ 0 & -A_{0,3}\kappa_1^3 & 2A_{1,2}\kappa_1^2\kappa_2 & -A_{2,1}\kappa_1\kappa_2^2 \\ 0 & A_{1,2}\kappa_1^2\kappa_2 & -2A_{2,1}\kappa_1\kappa_2^2 & A_{3,0}\kappa_2^3 \end{vmatrix} = (\kappa_1\kappa_2)^6\Delta.$$

Then  $\bar{D} = P_1 \cup P_2 \cup D$  where  $P_i$  denote parabolic line defined by  $\kappa_i = 0$ . We expect the behavior of the flow could be very degenerate along  $P_1$  and  $P_2$ , since the discriminant  $\bar{\Delta}$  is divisible by  $(\kappa_1\kappa_2)^6$

We assume that the rank of the matrix in (5.10) is 5. To investigate the equation  $\bar{C}(du, dv) = 0$ , we consider the foliation defined by the restriction of the 1-form  $\bar{\omega} = p\kappa_1du + q\kappa_2dv$  to  $M$ . The image of a flow of  $\bar{\omega}|_M$  by the projection  $\pi$  is a flow of  $\bar{C}(du, dv) = 0$ .

*Proof of Theorem 5.1.* We show the assertion at a point in  $M \cap \{p \neq 0\}$ . The case at a point in  $M \cap \{q \neq 0\}$  is similar and we omit the details. The equation  $C(1, \eta) = 0$  is an affine equation for  $M \cap \{p \neq 0\}$  where  $\eta = q/p$ , and this defines

- a function  $u$  of  $(v, \eta)$  by the implicit function theorem when  $C_u(1, \eta) \neq 0$ , and we have

$$u_v = -C_v(1, \eta)/C_u(1, \eta), \quad u_\eta = -C_q(1, \eta)/C_u(1, \eta).$$

- a function  $v$  of  $(u, \eta)$  by the implicit function theorem when  $C_v(1, \eta) \neq 0$ , and we have

$$v_u = -C_u(1, \eta)/C_v(1, \eta), \quad v_\eta = -C_q(1, \eta)/C_v(1, \eta).$$

Since  $C_u(\xi, 1)du + C_v(\xi, 1)dv + C_p(\xi, 1)d\xi = 0$ , we have

$$\begin{aligned} \frac{\bar{\omega}}{p} &= \kappa_1 du + \eta \kappa_2 dv \\ &= \begin{cases} \frac{1}{C_u(1, \eta)} [(C_u(1, \eta)\eta \kappa_2 - C_v(1, \eta)\kappa_1)dv - C_q(1, \eta)\kappa_1 d\eta] & C_u(1, \eta) \neq 0 \\ \frac{1}{C_v(1, \eta)} [(C_v(1, \eta)\kappa_1 - C_u(1, \eta)\eta \kappa_2)du - C_q(1, \eta)\eta \kappa_2 d\eta] & C_v(1, \eta) \neq 0 \end{cases} \end{aligned} \quad (5.16)$$

So if  $(p_0, q_0)$  is a multiple root of  $C(p, q) = 0$  at  $(u, v) = (0, 0)$  (i.e.,  $C_p(p_0, q_0) = C_q(p_0, q_0) = 0$  at  $(u, v) = (0, 0)$ ),  $\bar{\omega}|_M$  is singular at  $(0, 0) \times (p_0, q_0)$  if and only if  $\alpha q_0 \kappa_2 - \beta p_0 \kappa_1 = 0$ . So we obtain the condition 2. If  $(p_0, q_0)$  is not a multiple root of  $C(p, q) = 0$ , we have  $C_q(p_0, q_0) \neq 0$ . Now it is easy to conclude the condition 1.  $\square$

Let us consider the case that  $(0, 0)$  is parabolic (say  $k_1 = 0$ ). Then the tangent of the flow tend to the direction generated by  $\partial_u$ . Remark that  $\partial_u$  is not tangent to the parabolic line  $P_1$  whenever  $a_{30} \neq 0$ . If one writes the flow by  $t \mapsto (t, ct^2 + \dots)$ , we obtain that

$$a_{30}(k_2^3 c^3 - 3k_2^2 a_{21} c^2 + 3k_2 a_{12} a_{30} c - a_{03} a_{30}^2) = 0.$$

Observe that this has a multiple root when  $(0, 0) \in D$ .

**Proposition 5.7.** *Consider the case  $k_1 = 0$  and  $k_2 \neq 0$ . Assume that  $(0, 0)$  is ridge with respect to  $\mathbf{v}_1$  (i.e.,  $a_{30} = 0$ ). We assume that  $(0, 0)$  is not on the discriminant (i.e.,  $\Delta_0 = -a_{21}^2(3a_{12}^2 - 4a_{03}a_{21}) \neq 0$ ), For a single root  $(p_0, q_0)$  of  $c_3(p, q) = 0$ , we have the following.*

- If  $3a_{21}^2 < k_2 a_{40}$ , then  $\bar{\omega}|_M$  has a saddle at  $(0, 0) \times [p_0 : q_0]$ .
- If  $3a_{21}^2 > k_2 a_{40}$ , and  $2(k_2 a_{40}/2 - 3a_{21}^2)^2 < (k_2 a_{40})^2$ , then  $\bar{\omega}|_M$  has a node at  $(0, 0) \times [p_0 : q_0]$ .
- If  $2(k_2 a_{40}/2 - 3a_{21}^2)^2 > (k_2 a_{40})^2$ , then  $\bar{\omega}|_M$  has a focus at  $(0, 0) \times [p_0 : q_0]$ .

*Proof.* Suppose that  $p \neq 0$ ,  $C_u(p_0, q_0) \neq 0$ . Then  $u_v(0, 0) = -\beta/\alpha$ ,  $u_\eta(0, 0) = -C_q(1, \eta)/\alpha$ . We then observe

$$\begin{aligned} C_u(1, \eta)\eta \kappa_2 - C_v(1, \eta)\kappa_1 &= \frac{a_{21}}{k_2}(3a_{21}a_{12} - k_2 a_{31})v + (k_2 a_{40} - 3a_{21}^2)\eta + o(1) \\ -C_q(1, \eta)\kappa_1 &= -3a_{21}^2 v + o(1). \end{aligned}$$

The assertions are now concluded using the Lemma below, remarking the following:

$$\begin{aligned} \delta &= 3a_{21}^2(3a_{21}^2 - k_2 a_{40}), \quad \tau = k_2 a_{40}/2, \\ \delta - \tau^2 &= 3a_{21}^2(3a_{21}^2 - k_2 a_{40}) - k_2^2 a_{40}^2/4 = (k_2 a_{40}/2 - 3a_{21}^2)^2 - (k_2 a_{40})^2/2. \end{aligned}$$

$\square$


**Lemma 5.8.** Consider 1-form  $\omega = a(x, y)dx + b(x, y)dy$  on  $xy$  plane. Assume that  $a(0, 0) = b(0, 0) = 0$ . Set  $\delta = (a_x b_y - b_x a_y)(0, 0)$  and  $\tau = ((a_y - b_x)/2)(0, 0)$ . Then the singularity type of  $\omega$  at  $(0, 0)$  is saddle (resp. node, focus) if  $\delta < 0$  (resp.  $0 < \delta < \tau^2$ ,  $\tau^2 < \delta$ ).

*Proof.* It is enough to consider the singularity type of  $-b(x, y)\partial_x + a(x, y)\partial_y$ . We just compute the eigenvalue of the matrix

$$\begin{pmatrix} -b_x & -b_y \\ a_x & a_y \end{pmatrix} (0, 0)$$

and we have the result. □

We show below phase portraits of flows in  $uv$  plane.



**Remark 5.9.** We need to look the intersection of the parabolic line with the discriminant, since this can be considered as a degenerate point of cylindrical directions on the surface. We assume that  $(0, 0) \in P_1 \cap D$ , that is,  $k_1 = 0$  and

$$a_{30} = q_0^2 q_1, \quad a_{21} = -\frac{1}{3} q_0 (p_1 q_0 + 2p_0 q_1), \quad a_{12} = \frac{1}{3} p_0 (p_0 q_1 + 2p_1 q_0), \quad a_{03} = -p_0^2 p_1.$$

Whenever  $C_u(1, \eta_0) \neq 0$ ,  $u$  is a function of  $(v, \eta)$  by  $C(1, \eta) = A_{30} + 3A_{21}\eta + 3A_{12}\eta^2 + A_{03}\eta^3 = 0$ . By (5.16), we have

$$\begin{aligned} C_u(1, \eta) \frac{\bar{\omega}}{p} &= [C_u(1, \eta)\eta\kappa_2 - C_v(1, \eta)\kappa_1]dv - C_q(1, \eta)\kappa_1 d\eta \\ &= C_u(1, \eta)\eta\kappa_2 dv + o(1). \end{aligned}$$


A solution curve tending  $(0, 0)$  is expressed as

$$t \mapsto \left( t, -\frac{p_1 q_0^2}{2k_2} t^2 + \dots \right),$$

or, setting  $\epsilon = \text{sign} \frac{c_{4,u}(p_0, q_0)}{p_0 q_1 - p_1 q_0}$ ,

$$t \mapsto \left( -\epsilon t^2, -\frac{p_0 q_0 q_1}{2k_2} t^4 + \frac{2q_1}{5k_2} \sqrt{\left| \frac{c_{4,u}(p_0, q_0)}{p_0 q_1 - p_1 q_0} \right|} t^5 + \dots \right).$$

A phase portrait is given as follows.



**Remark 5.10.** Using Lemma 5.8, it should be possible to determine singularity types of  $\bar{\omega}|_M$  at points  $(0, 0) \times [p_0 : q_0]$  where  $[p_0 : q_0]$  are multiple roots of  $C(p, q) = 0$ .

Consider the equation  $C_q(1, \eta_0) = 0$ ,  $\eta_0 = q_0/p_0$ . We have

$$\begin{aligned} & \partial_v(C_u(1, \eta)\eta\kappa_2 - C_v(1, \eta)\kappa_1) \\ &= (- (C_v/C_u)C_{uu}(1, \eta) + C_{uv}(1, \eta))\eta\kappa_2 + C_u(1, \eta)\eta((-C_v/C_u)\kappa_{2,u} + \kappa_{2,v}) \\ & \quad - (- (C_v/C_u)C_{uv}(1, \eta) + C_{vv}(1, \eta))\kappa_1 - C_v(1, \eta)((-C_v/C_u)\kappa_{1,u} + \kappa_{1,v}) \\ &= (- (\beta/\alpha)C_{uu} + C_{uv})(1, \eta_0)\eta_0 k_2 - C_u(1, \eta_0)\eta_0(-a_{12}\beta/\alpha + a_{03}) \\ & \quad - (-C_{uv}\beta/\alpha + C_{vv})(1, \eta_0)k_1 - \beta((-\beta/\alpha)a_{30} + a_{21}) + o(0), \\ & \partial_\eta(C_u(1, \eta)\eta\kappa_2 - C_v(1, \eta)\kappa_1) \\ &= C_u(1, \eta)\kappa_2 + (- (C_q/C_u)C_{u,u}(1, \eta) + C_{uq}(1, \eta))\eta\kappa_2 - C_u(1, \eta)\eta(C_q/C_u)\kappa_{2,u} \\ & \quad - (- (C_q/C_u)C_{uv}(1, \eta) + C_{vq}(1, \eta))\kappa_1 - C_v(1, \eta)(C_q/C_u)\kappa_{1,u} \\ &= \alpha k_2 + C_{uq}(1, \eta_0)\eta_0 k_2 + \alpha \eta_0 a_{03} - C_{vq}(1, \eta_0)k_1 + o(0), \end{aligned}$$

$$\begin{aligned} \partial_v(-C_q(1, \eta)\kappa_1) &= - [-(C_v/C_u)C_{uq}(1, \eta_0) + C_{vq}(1, \eta_0)]\kappa_1 + C_q(1, \eta_0)\partial_v\kappa_1 \\ &= - (- (\beta/\alpha)C_{uq} + C_{vq})(1, \eta_0)k_1 + o(0), \\ \partial_\eta(-C_q(1, \eta)\kappa_1) &= - (- (C_q/C_u)C_{uq} + C_{qq})(1, \eta_0)\kappa_1 + C_q(1, \eta_0)\partial_\eta\kappa_1 \\ &= - C_{qq}(1, \eta_0)k_1 + o(0). \end{aligned}$$

So we obtain that

$$\delta = - [(- (\beta/\alpha)C_{uu} + C_{uv})(1, \eta_0)\eta_0 k_2 - C_u(1, \eta_0)\eta_0(-a_{12}\beta/\alpha + a_{03})]$$

$$\begin{aligned}
& -(-C_{uv}\beta/\alpha + C_{vv})(1, \eta_0)k_1 - \beta((-\beta/\alpha)a_{30} + a_{21})]C_{qq}(1, \eta_0)k_1 + C_{vq}(1, \eta_0)k_1 \\
& + [(-\beta/\alpha)C_{uq} + C_{vq})(1, \eta_0)k_1\alpha k_2 + C_{uq}(1, \eta_0)\eta_0 k_2 + \alpha\eta_0 a_{03}](-\beta/\alpha)C_{uq}, \\
\tau = & (\alpha + C_{uq}(1, \eta_0))k_2 + \alpha\eta_0 a_{03} - (\beta/\alpha)C_{uq}(1, \eta_0)k_1.
\end{aligned}$$

Now we can apply Lemma 5.8 to have a criterion for singularity types of  $\bar{\omega}|_M$ . But writing down the explicit conditions would be very complicated unfortunately. It could be better to discuss the normal form of jet and we discuss this method in the next subsection, which works in the generic context of cubic binary differential equations.

## 5.4 Reduction of jets of cubic differential equations

Now we discuss the reduction of jet of cubic binary differential equation  $\widehat{C}(du, dv) = 0$  where

$$\widehat{C}(p, q) = P(u, v)p^3 + 3Q(u, v)p^2q + 3R(u, v)pq^2 + S(u, v)q^3$$

where  $P = \sum_{i,j} \frac{p_{ij}}{i!j!} u^i v^j$ ,  $Q = \sum_{i,j} \frac{q_{ij}}{i!j!} u^i v^j$ ,  $R = \sum_{i,j} \frac{r_{ij}}{i!j!} u^i v^j$ ,  $S = \sum_{i,j} \frac{s_{ij}}{i!j!} u^i v^j$ . Set

$$\Delta_0 = \Delta(0, 0), \quad \Delta = \begin{vmatrix} P & 2Q & R & 0 \\ Q & 2R & S & 0 \\ 0 & P & 2Q & R \\ 0 & Q & 2R & S \end{vmatrix}.$$

**Proposition 5.11.**

1. If  $\Delta_0 > 0$  (resp.  $\Delta_0 < 0$ ), then the equation  $\widehat{C}(du, dv) = 0$  reduces to  $dx(dx^2 + 3dy^2) + o(0) = 0$  (resp.  $dx(dx^2 - 3dy^2) + o(0) = 0$ ).
2. When  $\Delta_0 = 0$ , then the constant term of  $\widehat{C}(du, dv)$  is  $(q_0 du - p_0 dv)^2(q_1 du - p_1 dv)$ . Now we set

$$\sigma_0 = \alpha p_0 + \beta q_0, \quad \sigma_1 = \alpha p_1 + \beta q_1, \quad \sigma_2 = p_0 q_1 - p_1 q_0,$$

where  $\alpha = \widehat{C}_u(p_0, q_0)|_{(u,v)=(0,0)}$  and  $\beta = \widehat{C}_v(p_0, q_0)|_{(u,v)=(0,0)}$ .

- (a) When  $\sigma_0 \neq 0$ ,  $\sigma_1 \neq 0$ , and  $\sigma_2 \neq 0$ , the equation reduces to

$$[dx^2 + (x + y)dy^2]dy + o(1) = 0. \quad (5.17)$$

- (b) When  $\sigma_0 \neq 0$ ,  $\sigma_1 = 0$ , and  $\sigma_2 \neq 0$ , the equation reduces to

$$[dx^2 + (ax^2 + y)dy^2]dy + o(2) = 0. \quad (5.18)$$

When  $a \neq 0$ , the discriminant has the first order tangent to the foliation corresponding to the single root of the cylindrical equation. The sign of  $a$  coincides with the sign of (5.22) in the proof.

(c) When  $\sigma_0 = 0$ ,  $\sigma_1 \neq 0$ , and  $\sigma_2 \neq 0$ , the equation reduces to

$$[dx^2 + (x + by^2)dy^2]dy + o(2) = 0. \quad (5.19)$$

The equation (5.19) defines folded saddle (resp. folded node, folded focus) if  $b < 0$  (resp.  $0 < b < 1/16$ ,  $b > 1/16$ ). Remark that  $b$  is given by (5.23).

3. If  $\Delta_0 = \sigma_2 = 0$ , then the equation reduces to

$$dx^3 + 3(kx + ly)dx dy^2 + (ax + by)dy^3 + o(1) = 0 \quad (5.20)$$

and the singularity type of the discriminant is cusp if  $al - bk \neq 0$ , or equivalently

$$\begin{vmatrix} \widehat{C}_{up}(p_0, q_0) & \widehat{C}_{vp}(p_0, q_0) \\ \widehat{C}_{uq}(p_0, q_0) & \widehat{C}_{vq}(p_0, q_0) \end{vmatrix} \neq 0. \quad (5.21)$$

As we will see in the proof, explicit formulas for  $a$  in (5.18) and for  $b$  in (5.19) are very complicate when we express them in terms of the coefficients of Monge normal form (2.1).

*Proof.* By a suitable linear change of coordinates, the 0-jet of  $\widehat{C}$  reduces to  $dx(dx^2 + 3dy^2)$  (resp.  $dx(dx^2 - 3dy^2)$ ) when  $\Delta_0 > 0$  (resp.  $\Delta_0 < 0$ ). If this determinant is 0, we can assume that the 0-jet of the equation is  $(q_0 du - p_0 dv)^2(q_1 du - p_1 dv)$  and we reduce the 0-jet to  $3\sigma_2^3 t_0^2 t_1 dx^2 dy$  by a coordinates change given by

$$u = t_1 p_1 x + t_0 p_0 y + o(1), \quad v = t_1 q_1 x + t_0 q_0 y + o(1),$$

whenever  $\sigma_2 \neq 0$ . Here  $t_0, t_1$  are non-zero constants. Multiplying the equation by  $1/(3\sigma_2^3 t_0^2 t_1)$ , we reduce the constant term of the coefficient of  $dx^2 dy$  is 1. In this case, by a suitable choice of quadratic parts of coordinate change, we are able to reduce the equation to

$$[dx^2 + (a_1 x + a_2 y)dy^2]dy + o(1) = 0$$

where  $a_1 = t_0^2 \sigma_1 / (3t_1 \sigma_2^3)$ ,  $a_2 = t_0^3 \sigma_0 / (3t_1^2 \sigma_2^3)$ . When  $\sigma_0 \neq 0$ ,  $\sigma_1 \neq 0$ , we reduce this equation to (5.17) setting  $(t_0, t_1) = (3\sigma_0 \sigma_2^5 / \sigma_1^3)(1, \sigma_0 \sigma_2^2)$ . Remark also that, if  $a_1 \neq 0$ , then we reduce the coefficient of  $x dy^3$  to 1 setting  $t_1 = t_0^2 \sigma_1 / (3\sigma_2^3)$ .

When  $\sigma_0 \neq 0$ ,  $\sigma_1 = 0$  (i.e.,  $a_1 = 0$ ,  $a_2 \neq 0$ ), we need to look the coefficients of  $x^2 dy^3$  which is expressed as follows:

$$t_0^2 \left[ \frac{3(\widehat{C}_{uu}(p_0, q_0)p_1^2 + 2\widehat{C}_{uv}(p_0, q_0)p_1 q_1 + \widehat{C}_{vv}(p_0, q_0)q_1^2)}{4\sigma_2^3} - \frac{\widehat{Q}}{12\sigma_2^6} \right] \quad (5.22)$$

where  $\widehat{Q} = \sum_{i,j} Q_{ij} p_0^{4-j} q_0^j p_1^{4-i} q_1^i$ ,  $Q_{00} = 25p_{10}^2$ ,

$$Q_{01} = 37p_{01}p_{10} + 63p_{10}q_{10}, \quad Q_{02} = 12p_{01}^2 + 30p_{10}q_{01} + 57p_{01}q_{10} + 9q_{10}^2 + 42p_{10}r_{10},$$

$$\begin{aligned}
Q_{03} &= 24p_{01}q_{01} + 18q_{01}q_{10} + 9p_{10}r_{01} + 36p_{01}r_{10} + 13p_{10}s_{10}, \\
Q_{04} &= 9q_{01}^2 + 3p_{01}r_{01} + 2p_{10}s_{01} + 11p_{01}s_{10}, \quad Q_{10} = 13p_{01}p_{10} + 87p_{10}q_{10}, \\
Q_{11} &= 13p_{01}^2 + 114p_{10}q_{01} + 60p_{01}q_{10} + 171q_{10}^2 + 42p_{10}r_{10}, \\
Q_{12} &= 87p_{01}q_{01} + 225q_{01}q_{10} + 75p_{10}r_{01} + 48p_{01}r_{10} + 162q_{10}r_{10} + 3p_{10}s_{10}, \\
Q_{13} &= 54q_{01}^2 + 81p_{01}r_{01} + 63q_{10}r_{01} + 144q_{01}r_{10} + 14p_{10}s_{01} + 5p_{01}s_{10} + 39q_{10}s_{10}, \\
Q_{14} &= 45q_{01}r_{01} + 16p_{01}s_{01} + 6q_{10}s_{01} + 33q_{01}s_{10}, \\
Q_{20} &= 6p_{10}q_{01} + 33p_{01}q_{10} + 45q_{10}^2 + 66p_{10}r_{10}, \\
Q_{21} &= 39p_{01}q_{01} + 207q_{01}q_{10} + 60p_{10}r_{01} + 33p_{01}r_{10} + 243q_{10}r_{10} + 18p_{10}s_{10}, \\
Q_{22} &= 162q_{01}^2 + 27p_{01}r_{01} + 324q_{10}r_{01} + 162q_{01}r_{10} + 162r_{10}^2 + 18p_{10}s_{01} + 18p_{01}s_{10} + 27q_{10}s_{10}, \\
Q_{23} &= 243q_{01}r_{01} + 207r_{01}r_{10} + 18p_{01}s_{01} + 60q_{10}s_{01} + 33q_{01}s_{10} + 39r_{10}s_{10}, \\
Q_{24} &= 45r_{01}^2 + 66q_{01}s_{01} + 6r_{10}s_{01} + 33r_{01}s_{10}, \\
Q_{30} &= 6p_{10}r_{01} + 33p_{01}r_{10} + 45q_{10}r_{10} + 16p_{10}s_{10}, \\
Q_{31} &= 39p_{01}r_{01} + 63q_{10}r_{01} + 144q_{01}r_{10} + 54r_{10}^2 + 14p_{10}s_{01} + 5p_{01}s_{10} + 81q_{10}s_{10}, \\
Q_{32} &= 162q_{01}r_{01} + 225r_{01}r_{10} + 3p_{01}s_{01} + 75q_{10}s_{01} + 48q_{01}s_{10} + 87r_{10}s_{10}, \\
Q_{33} &= 171r_{01}^2 + 42q_{01}s_{01} + 114r_{10}s_{01} + 60r_{01}s_{10} + 13s_{10}^2, \quad Q_{34} = 87r_{01}s_{01} + 13s_{01}s_{10}, \\
Q_{40} &= 9r_{10}^2 + 2p_{10}s_{01} + 11p_{01}s_{10} + 3q_{10}s_{10}, \\
Q_{41} &= 18r_{01}r_{10} + 13p_{01}s_{01} + 9q_{10}s_{01} + 36q_{01}s_{10} + 24r_{10}s_{10}, \quad Q_{44} = 25s_{01}^2, \\
Q_{42} &= 9r_{01}^2 + 42q_{01}s_{01} + 30r_{10}s_{01} + 57r_{01}s_{10} + 12s_{10}^2, \quad Q_{43} = 63r_{01}s_{01} + 37s_{01}s_{10}.
\end{aligned}$$

By a suitable choice of cubic parts of the coordinate change we are able to reduce the equation to (5.18). Remark that this coordinate change does not change the coefficient of  $x^2dy^3$ . We also remark that the possible singularity of  $\omega|_M$  over  $(0, 0)$  is  $[p_0, q_0] = [0 : 1]$  only if  $a_2 = 0$ .

We consider the case  $\sigma_0 = 0$  and  $\sigma_1 \neq 0$  (i.e.,  $a_1 \neq 0$ ,  $a_2 = 0$ ). We need to look the coefficients of  $y^2dy^3$  which is expressed as follows:

$$\left( \frac{t_0}{t_1} \right)^2 \left[ \frac{\widehat{C}_{uu}(p_0, q_0)p_0^2 + 2\widehat{C}_{uv}(p_0, q_0)p_0q_0 + \widehat{C}_{vv}(p_0, q_0)q_0^2}{6\sigma_2^3} - \frac{\widehat{R}}{4\sigma_2^6} \right]$$

where  $\widehat{R} = \sum_{i,j} R_{ij} p_0^{6-j} q_0^j p_1^{2-i} q_1^i$ ,

$$\begin{aligned}
R_{00} &= 4p_{10}^2, \quad R_{01} = 3p_{10}(7p_{01} + q_{10}), \quad R_{02} = 17p_{01}^2 + 28p_{10}q_{01} + 43p_{01}q_{10} - 24q_{10}^2 - 4p_{10}r_{10}, \\
R_{03} &= 68p_{01}q_{01} + 30q_{01}q_{10} + 7p_{10}r_{01} + 37p_{01}r_{10} - 63q_{10}r_{10} + p_{10}s_{10}, \\
R_{04} &= 3(18q_{01}^2 + 16p_{01}r_{01} - 9q_{10}r_{01} + 6q_{01}r_{10} - 12r_{10}^2 + 5p_{01}s_{10} - 4q_{10}s_{10}), \\
R_{05} &= 54q_{01}r_{01} - 33r_{01}r_{10} + 14p_{01}s_{01} - 14q_{10}s_{01} + 16q_{01}s_{10} - 13r_{10}s_{10}, \\
R_{06} &= 3r_{01}^2 + 14q_{01}s_{01} - 14r_{10}s_{01} + r_{01}s_{10}, \\
R_{10} &= -p_{10}(13p_{01} - 21q_{10}), \quad R_{11} = -13p_{01}^2 + 10p_{10}q_{01} - 20p_{01}q_{10} + 57q_{10}^2 + 14p_{10}r_{10}, \\
R_{12} &= -31p_{01}q_{01} + 75q_{01}q_{10} + 31p_{10}r_{01} - 29p_{01}r_{10} + 81q_{10}r_{10} - 7p_{10}s_{10}, \\
R_{13} &= 2(9q_{01}^2 - 6p_{01}r_{01} + 63q_{10}r_{01} + 18q_{01}r_{10} + 9r_{10}^2 + 4p_{10}s_{01} - 11p_{01}s_{10} - 6q_{10}s_{10}),
\end{aligned}$$

$$\begin{aligned}
R_{14} &= 81q_{01}r_{01} + 75r_{01}r_{10} - 7p_{01}s_{01} + 31q_{10}s_{01} - 29q_{01}s_{10} - 31r_{10}s_{10}, \\
R_{15} &= 57r_{01}^2 + 14q_{01}s_{01} + 10r_{10}s_{01} - 20r_{01}s_{10} - 13s_{10}^2, \quad R_{16} = s_{01}(21r_{01} - 13s_{10}), \\
R_{20} &= -14p_{10}q_{01} + p_{01}q_{10} + 3q_{10}^2 + 14p_{10}r_{10}, \\
R_{21} &= -13p_{01}q_{01} - 33q_{01}q_{10} - 14p_{10}r_{01} + 16p_{01}r_{10} + 54q_{10}r_{10} + 14p_{10}s_{10}, \\
R_{22} &= -3(12q_{01}^2 + 4p_{01}r_{01} + 9q_{10}r_{01} - 6q_{01}r_{10} - 18r_{10}^2 - 5p_{01}s_{10} - 16q_{10}s_{10}), \\
R_{23} &= -63q_{01}r_{01} + 30r_{01}r_{10} + p_{01}s_{01} + 7q_{10}s_{01} + 37q_{01}s_{10} + 68r_{10}s_{10}, \\
R_{24} &= 28r_{10}s_{01} + 43r_{01}s_{10} + 17s_{10}^2 - 24r_{01}^2 - 4q_{01}s_{01}, \quad R_{25} = 3s_{01}(r_{01} + 7s_{10}), \quad R_{26} = 4s_{01}^2.
\end{aligned}$$

We assume that  $a_1 = 1$ , since we can choose  $(t_0, t_1)$  with  $t_0^2/t_1 = 3\sigma_2^3/\sigma_1$ . Then the coefficient of  $y^2dy^3$  becomes

$$\frac{9}{\sigma_1^2} \left[ \frac{\sigma_2^3(\widehat{C}_{uu}(p_0, q_0)p_0^2 + 2\widehat{C}_{uv}(p_0, q_0)p_0q_0 + \widehat{C}_{vv}(p_0, q_0)q_0^2)}{6} - \frac{\widehat{R}}{4} \right]. \quad (5.23)$$

By a suitable choice of cubic parts of the coordinate change, we are able to reduce the equation to (5.19). Remark that this coordinate change does not change the coefficient of  $y^2dy^3$ . Now we can apply Lemma 5.8 to have a criterion for singularity types of the restriction of 1-form  $pdu - qdv$  to the subset  $\widehat{C}(p, q) = 0$  in  $(\mathbb{R}^2, 0) \times P^1$ .

Remark that the linear term of  $\Delta$  is  $\frac{4}{27}\sigma_2^3(a_1x + a_2y)$  and  $(0, 0) \in \Sigma D$  if and only if  $\sigma_2 = 0$  or  $a_1 = a_2 = 0$ .

In the case  $\sigma_2 = 0$ , the constant part of  $\widehat{C}(du, dv)$  is a constant multiple of  $(v_0du - u_0dv)^3$  and we are able to reduce the 0-jet of the equation to  $dx^3$ . By a suitable choice of quadratic parts of coordinate change, we are able to reduce the equation to (5.19). We remark that


$$\begin{aligned}
k &= -p_0^{-3}t^{-1}\widehat{C}_{vq}(p_0, q_0), & l &= -p_0^{-3}t^{-2}[\widehat{C}_{uq}(p_0, q_0)p_0 + \widehat{C}_{vq}(p_0, q_0)q_0], \\
a &= -p_0^{-3}t^{-2}\widehat{C}_v(p_0, q_0), & b &= -p_0^{-3}t^{-3}[\widehat{C}_u(p_0, q_0)p_0 + \widehat{C}_v(p_0, q_0)q_0] \\
(\text{or } k &= q_0^{-3}t^{-1}\widehat{C}_{uq}(p_0, q_0), & l &= q_0^{-3}t^{-2}[\widehat{C}_{uq}(p_0, q_0)p_0 + \widehat{C}_{vq}(p_0, q_0)q_0], \\
a &= q_0^{-3}t^{-2}\widehat{C}_u(p_0, q_0), & b &= q_0^{-3}t^{-3}[\widehat{C}_u(p_0, q_0)p_0 + \widehat{C}_v(p_0, q_0)q_0]),
\end{aligned}$$


by a suitable change of coordinate

$$u = p_0y + o(1), \quad v = tx + q_0y + o(1) \quad (\text{or } u = tx + p_0y + o(1), \quad v = q_0y + o(1))$$

and the 3-jet of the discriminant is  $(ax + by)^2 + (kx + ly)^3$ . So the singularity type of the discriminant is cusp if  $al - bk \neq 0$ , which is equivalent to (5.21).  $\square$

The phase portraits of singularities of the flows are shown by the pictures below.





We can apply Proposition 5.11 to determine the singularity type of the equation

$$\bar{C}(du, dv) = 0,$$

whenever  $k_1 k_2 \neq 0$ .

## References

- [1] T. Banchoff, T. Gaffney, and C. McCrory, *Cusps of Gauss maps*, Research Notes in Mathematics 55, Pitman, 1982.
- [2] J. W. Bruce and P. J. Giblin, *Curves and Singularities*, Cambridge University Press, 1984. Second Edition, 1992.
- [3] J. W. Bruce, P. J. Giblin, and F. Tari, Families of surfaces: height functions, Gauss maps and duals, in “Real and Complex Singularities”, Ed. W. L. Marar, Pitman Research Notes in Math. 333 (1995).
- [4] J. W. Bruce and F. Tari, Dupin indicatrices and families of curve congruences, *Trans. Amer. Math. Soc.* 357 (2005), 267–285.
- [5] G. Darboux, *Leçons sur la théorie générale des surfaces*, IV, Gauthier-Villars, Paris, 1896.
- [6] T. Fukui and M. Hasegawa, Singularities of parallel surfaces, *Tohoku Math. J.* (2) 64 (2012), no. 3, 387–408. 2015/1/30
- [7] S. Izumiya, Differential geometry from the viewpoint of Lagrangian or Legendrian singularity theory, in *Singularity theory, Proceedings of the Singularity school and conference, Marceille, 2005* (D. Chéniot, N. Dutertre, C. Murolo, D. Trotman and A. Pichon) World Scientific Pub Co Inc 2007.

- [8] S. Izumiya, M. C. Romero Fuster, M. A. Ruas, and F. Tari, Differential geometry from singularity theory viewpoint, World Scientific Pub Co Inc (2015/1/30).
- [9] D. K. H. Mochida, M. C. Romero Fuster, and M. A. Ruas, The geometry of surfaces in 4-space from a contact viewpoint. Geometriae Dedicata 54 (1995), 323–332.
- [10] R. A. Garcia, D. K. H. Mochida, M. C. Romero Fuster and M. A. Ruas, Inflection points and topology of surfaces in 4-space, Trans. Amer. Math. Soc. 352 (2000), 3029–3043.
- [11] K. Kakié, The resultant of several homogeneous polynomials in two indeterminates, Proc. Amer. Math. Soc. 54 (1976), 1–7.
- [12] J. J. Koenderink, What does the occluding contour tell us about solid shape?, Perception, 13 (1984), 321–330.
- [13] J. J. Koenderink, Solid shape, MIT Press Series in Artificial Intelligence, MIT Press, Cambridge, MA, 1990.
- [14] J. Mather, Stability of mappings, III. Finitely determined map-germs, Publ. Math. I.H.E.S. 35(1969), 127–156.
- [15] J. Montaldi, On contact between submanifolds, Michigan Math. J. 33 (1986), 195–199.
- [16] I. R. Porteous, The normal singularities of a submanifold, J. Differential Geometry 5 (1971), 543–564.
- [17] I. R. Porteous, The normal singularities of surfaces in  $\mathbb{R}^3$ , Singularities, Part 2 (Arcata, Calif., 1981), 379–393, Proc. Sympos. Pure Math., 40, Amer. Math. Soc., Providence, RI, 1983.
- [18] T. Takahashi, Homogeneous hypersurfaces in space of constant curvature, J. Math. Soc. Japan 22, (1970), 395–410.
- [19] R. Thom, Stabilité structurelle et morphogénèse, Benjamin Paris, 1972.

Toshizumi Fukui,

Department of Mathematics, Faculty of Science, Saitama University, Saitama, 338-8570, Japan

*E-mail address:* `tfukui@rimath.saitama-u.ac.jp`

Masaru Hasegawa,

Departamento de Matemática, Instituto de Ciências Matemáticas e de Computação, Universidade de São Paulo - Campus de São Carlos, Caixa Postal 668, 13560-970 São Carlos, SP, Brazil

*E-mail address:* `mhasegawa@icmc.sc.usp.br`

Kouichi Nakagawa,

Department of Mathematics, Faculty of Science, Saitama University, Saitama, 338-8570, Japan