

On the anti-canonical ring and varieties of Fano type

Paolo Cascini and Yoshinori Gongyo

Dedicated to Professor Fumio Sakai on the occasion of his retirement.

(Received 23 July, 2013; Revised 24 September, 2013; Accepted 1 October, 2013)

Abstract

We study the anti-canonical ring of a projective variety and we characterise varieties of log Fano type depending on the singularities of these models.

1. Introduction

Over the last few years, there have been many developments about the study of the canonical model of a projective variety. On the other hand, not much is known about its anti-canonical model. Few decades ago, Fumio Sakai considered this problem for projective surfaces (e.g. see [Sk2], and [Sk3]). In particular, he gave a characterisation of the anti-canonical model of any rational surface with big anti-canonical divisor.

The purpose of these notes is to study the anti-canonical model for higher dimensional projective varieties. More specifically, we describe a characterisation of varieties of Fano type depending on the singularities of their anti-canonical models.

Our main result is the following:

Theorem 1.1 (=Corollary 3.2). *Let X be a \mathbb{Q} -Gorenstein normal variety.*

Then X is of Fano type if and only if $-K_X$ is big, $R(-K_X)$ is finitely generated, and $\text{Proj } R(-K_X)$ has log terminal singularities.

Recall that a normal projective variety X is said to be of *Fano type* if it admits a \mathbb{Q} -divisor $\Delta \geq 0$ such that (X, Δ) is klt and $-(K_X + \Delta)$ is ample. In dimension two, the main result of [TVAV] implies that a rational surface with big anti-canonical divisor is a Mori dream space. Thus, the theorem follows from the arguments in [GOST]. On the other hand, it is known that higher dimensional varieties with big and nef anti-log canonical divisor are not necessarily Mori dream spaces (e.g. see [G, Example 5.2]). Note that we do not assume that X is a Mori dream space in Theorem 1.1. Note that in dimension two, Hwang and

2010 Mathematics Subject Classification. Primary 14J45, Secondary 14E30

Key words and phrases. Fano variety, anti-canonical ring, singularity of log pair

The first author was partially supported by an EPSRC grant. The second author was partially supported by Aid-in-Grant from JSPS and Research expense from the JRF fund.

Park gave a different proof of Theorem 1.1 (see [HP, Theorem 1.1])

The paper is organised as follows. We give a proof of Theorem 1.1 in Section 3. In Section 4, we consider the problem about the finite generation of Cox rings of log canonical log Fano varieties and in Section 5 we describe some examples of varieties of Fano type which give a negative answer to some questions raised before on these problems.

2. Preliminaries

We work over the field of complex numbers. We use the standard definitions for the singularities appearing in the minimal model program:

Definition 2.1 (cf. [KoM, Definition 2.34]). Let X be a normal variety and Δ be an effective \mathbb{Q} -divisor on X such that $K_X + \Delta$ is \mathbb{Q} -Cartier. Let $\pi : \tilde{X} \rightarrow X$ be a birational morphism from a normal variety \tilde{X} . Then we can write

$$K_{\tilde{X}} = \pi^*(K_X + \Delta) + \sum_E a(E, X, \Delta)E,$$

where E runs through all the distinct prime divisors on \tilde{X} and $a(E, X, \Delta)$ is a rational number. We say that the pair (X, Δ) is *log canonical* (resp. *klt*) if $a(E, X, \Delta) \geq -1$ (resp. $a(E, X, \Delta) > -1$) for every prime divisor E over X . If $\Delta = 0$, we simply say that X has only log canonical singularities (resp. log terminal singularities). We say that the pair (X, Δ) is *dlt* if the coefficients of Δ are less or equal one and $a(E, X, \Delta) > -1$ for any exceptional divisor E of π .

2.1 Section rings and Cox rings

Definition 2.2. Let X be a normal variety and let L_i be a line bundle for $i = 1, \dots, k$. We define the ring

$$R(X; L_1, \dots, L_k) = \sum_{(m_1, \dots, m_k) \in \mathbb{Z}} H^0(X, \otimes_{i=1}^k L_i^{m_i}).$$

In particular, if $\{L_i\}$ is a basis of $\text{Pic}_{\mathbb{Q}} X$, the ring $R(X; L_1, \dots, L_k)$ is called the *Cox ring* of X (see [GOST, Subsection 2.4] for more details).

2.2 Mori dream spaces

We now recall a few basic facts about Mori Dream Spaces, introduced by Hu and Keel in [HK] (see also [M]).

Definition 2.3. A normal projective variety X over a field is called a *\mathbb{Q} -factorial Mori dream space* (or *Mori dream space* for short) if X satisfies the following three conditions:

- (i) X is \mathbb{Q} -factorial, $\text{Pic}(X)$ is finitely generated, and $\text{Pic}(X)_{\mathbb{Q}} \simeq \mathbb{N}^1(X)_{\mathbb{Q}}$,
- (ii) $\text{Nef}(X)$ is the affine hull of finitely many semi-ample line bundles,
- (iii) there exists a finite collection of small birational maps $f_i : X \dashrightarrow X_i$ such that each X_i satisfies (i) and (ii), and that $\text{Mov}(X)$ is the union of the $f_i^*(\text{Nef}(X_i))$.

Remark 2.4. *Over the field of complex numbers, the finite generation of $\text{Pic}(X)$ is equivalent to the condition $\text{Pic}(X)_{\mathbb{Q}} \simeq \mathbb{N}^1(X)_{\mathbb{Q}}$.*

As suggested by the name, one of the advantages of working on a Mori dream space, is that on these varieties we can run an MMP for any divisor:

Proposition 2.5. *([HK, Proposition 1.11]) Let X be a \mathbb{Q} -factorial Mori dream space. Then for any divisor D on X which is not nef, there exists always a D -negative contraction or a flip. In addition, any sequence of such flips terminates.*

We will refer to such a sequence as a D -MMP.

The following known result is a useful source of Mori dream spaces:

Proposition 2.6 (cf. [Br, Theorem 3.2]). *Let X be a \mathbb{Q} -factorial projective variety which is a Mori dream space and let L_1, \dots, L_m be Cartier divisors on X . Let*

$$\mathcal{E} = \bigoplus_{i=1}^m \mathcal{O}_X(L_i)$$

and let $f : Y = \mathbb{P}(\mathcal{E}) \rightarrow X$ be the corresponding projective bundle.

Then Y is also a Mori dream space.

Proof. Let $\{H_i\}_{i=1}^k$ be a basis of $\text{Pic}_{\mathbb{Z}} X$. Then we can write

$$L_i = \sum a_{i,j} H_j,$$

where $a_{i,j}$ are integers. Let S be the tautological section of \mathcal{E} , i.e. the section corresponding to $\mathcal{O}_Y(1)$. Then for any $b_0, \dots, b_k \in \mathbb{Z}$, it follows that

$$\begin{aligned} & H^0(Y, b_0 S + \sum_j b_j f^* H_j) \\ & \simeq \bigoplus_{d_1 + \dots + d_m = b_0, d_j \geq 0} H^0(X, \sum_j d_j L_j + \sum_i b_i H_i) t_1^{d_1} \dots t_m^{d_m} \\ & = \bigoplus_{d_1 + \dots + d_m = b_0, d_j \geq 0} H^0(X, \sum_i (b_i + \sum_j d_j a_{j,i}) H_i) t_1^{d_1} \dots t_m^{d_m} \end{aligned}$$

when $b_0 \geq 0$ and

$$H^0(Y, b_0 S + \sum_{j=1}^k b_j f^* H_j) = 0$$

when $b_0 < 0$.

Thus $R(Y; S, f^* H_1, \dots, f^* H_k)$ is a direct summand of the ring

$$R(X; H_1, \dots, H_k)[t_1, \dots, t_m].$$

Since $R(Y; S, f^* H_1, \dots, f^* H_k)$ is finitely generated and

$$\{S, f^* H_1, \dots, f^* H_k\}$$

is a basis of $\text{Pic}_{\mathbb{Z}} Y$, we have that Y is also a Mori dream space by [HK, Proposition 2.9]. \square

2.3 Varieties of Fano type and of Calabi–Yau type

Definition 2.7 (cf. [PS, Lemma-Definition 2.6]). Let X be a projective normal variety and Δ be an effective \mathbb{Q} -divisor on X such that $K_X + \Delta$ is \mathbb{Q} -Cartier.

- (i) We say that (X, Δ) is a *log Fano pair* if $-(K_X + \Delta)$ is ample and (X, Δ) is klt. We say that X is of *Fano type* if there exists an effective \mathbb{Q} -divisor Δ on X such that (X, Δ) is a log Fano pair.
- (ii) We say that (X, Δ) is a *log Calabi–Yau pair* if $K_X + \Delta \sim_{\mathbb{Q}} 0$ and (X, Δ) is log canonical. We say that X is of *Calabi–Yau type* if there exists an effective \mathbb{Q} -divisor Δ such that (X, Δ) is log Calabi–Yau.

Remark 2.8. *If there exists an effective \mathbb{Q} -divisor Δ on X such that (X, Δ) is klt and $-(K_X + \Delta)$ is nef and big, then X is of Fano type. See [PS, Lemma-Definition 2.6].*

Similarly to Proposition 2.6, we have:

Proposition 2.9. *Let X be a projective variety of Fano type and let L_1, \dots, L_m be Cartier divisors on X . Let*

$$\mathcal{E} = \bigoplus_{i=1}^m \mathcal{O}_X(L_i)$$

and let $f: Y = \mathbb{P}(\mathcal{E}) \rightarrow X$ be the corresponding projective bundle.

Then Y is also of Fano type

Proof. By assumption, there exists a klt pair (X, Δ) such that $-(K_X + \Delta)$ is ample. We may assume that $\Delta = A + B$ where A is an ample \mathbb{Q} -divisor and (X, B) is klt.

After tensoring by a suitable ample divisor on X , we may assume that

L_1, \dots, L_m are very ample divisors on X . Let S be the tautological section of \mathcal{E} , i.e. the section corresponding to $\mathcal{O}_Y(1)$. Since L_1, \dots, L_m are ample, it follows that S is ample on Y .

Let T_1, \dots, T_m be the divisors in Y given by the summands of \mathcal{E} . Then

$$mS \sim \sum_{i=1}^m (T_i + f^*L_i).$$

Fix a sufficiently small rational number $\delta > 0$ such that $f^*A - \delta \sum_{i=1}^m f^*L_i$ is ample. Then for any sufficiently small rational number $\varepsilon > 0$, there exists a \mathbb{Q} -divisor $\Gamma \geq 0$ on Y such that (Y, Γ) is klt and

$$\begin{aligned} \sum_{i=1}^m T_i - \varepsilon S + f^*\Delta &= \sum_{i=1}^m T_i + f^*A - \varepsilon S + f^*B \\ &\sim_{\mathbb{Q}} (1 - \delta) \sum_{i=1}^m T_i + (m\delta - \varepsilon)S + f^*(A - \delta \sum_{i=1}^m L_i) + f^*B \\ &\sim_{\mathbb{Q}} \Gamma \end{aligned}$$

In addition, we have

$$K_Y = - \sum_{i=1}^m T_i + f^*K_X$$

and it follows that

$$-(K_Y + \Gamma) \sim_{\mathbb{Q}} -(K_Y + \sum_{i=1}^m T_i) - f^*\Delta + \varepsilon S = -f^*(K_X + \Delta) + \varepsilon S.$$

Thus, if ε is sufficiently small, then (Y, Γ) is log Fano and the claim follows. \square

Proposition 2.10. *Let X be a projective variety of Calabi–Yau type and let L_1, \dots, L_m be Cartier divisors on X . Let*

$$\mathcal{E} = \bigoplus_{i=1}^m \mathcal{O}_X(L_i)$$

and let $f: Y = \mathbb{P}(\mathcal{E}) \rightarrow X$ be the corresponding projective bundle.

Then Y is also of Calabi–Yau type

Proof. By assumption, there exists a log canonical pair (X, Δ) such that $K_X + \Delta \sim_{\mathbb{Q}} 0$.

Let T_1, \dots, T_m be the divisors in Y given by the summands of \mathcal{E} . Then there exist a \mathbb{Q} -divisor $\Gamma \geq 0$ on Y such that (Y, Γ) is log canonical and

$$\Gamma \sim_{\mathbb{Q}} \sum_{i=1}^m T_i + f^*\Delta.$$

In addition, we have

$$K_Y = - \sum_{i=1}^m T_i + f^* K_X$$

and it follows that

$$K_Y + \Gamma \sim_{\mathbb{Q}} f^*(K_X + \Delta).$$

Thus, (Y, Γ) is log Calabi–Yau and the claim follows. \square

3. The anti-canonical model of a variety of Fano type

We begin with a generalisation of Theorem 1.1:

Theorem 3.1. *Let (X, Δ) be a projective log pair over an algebraic closed field k of characteristic zero, such that $K_X + \Delta$ is \mathbb{Q} -Cartier.*

Then there exists a \mathbb{Q} -divisor $\Delta' \geq 0$ such that $(X, \Delta + \Delta')$ is a log Fano pair if and only if $R(X, -m(K_X + \Delta))$ is a finitely generated k -algebra for some $m \in \mathbb{N}$, the map

$$\varphi : X \dashrightarrow Y := \text{Proj } R(X, -m(K_X + \Delta))$$

is birational and the pair $(Y, \Gamma = \varphi_ \Delta)$ is klt.*

Proof. We first prove the "only if" part. To this end, we follow some of the arguments in [GOST]. Assume that the pair $(X, \Delta + \Delta')$ is log Fano, for some \mathbb{Q} -divisor $\Delta' \geq 0$. After possibly replacing X by its \mathbb{Q} -factorization [BCHM, Corollary 1.4.3], we may assume that X is \mathbb{Q} -factorial. In particular $(X, \Delta + \Delta')$ is klt and $A := -(K_X + \Delta + \Delta')$ is ample. Thus $R(X, -m(K_X + \Delta))$ is finitely generated for some positive integer m . In addition, $-(K_X + \Delta)$ is big and, in particular, φ is a birational map. Let D be an effective \mathbb{Q} -divisor such that $D \sim_{\mathbb{Q}} -(K_X + \Delta)$ and pick a rational number $0 < \varepsilon \ll 1$ such that $(X, \Delta + \Delta' + \varepsilon D)$ is klt. Then

$$\varepsilon D = K_X + \Delta + \Delta' + A + \varepsilon D$$

and by [BCHM], we can run a D -MMP and obtain a contraction map $\psi : X \dashrightarrow X'$ such that $\psi_* D$ is big and semi-ample. In particular $(X', \psi_* \Delta)$ is klt. Note that φ factors through ψ . Let $\eta : X' \rightarrow Y$ be the induced morphism. Then, $(Y, \Gamma = \eta_* \psi_* \Delta)$ is klt.

Now we show the "if" part. By assumption

$$\varphi : X \dashrightarrow Y := \text{Proj } R(X, -m(K_X + \Delta))$$

is a birational contraction and $-(K_Y + \Gamma) = -\varphi_*(K_X + \Delta)$ is \mathbb{Q} -Cartier by [HK, Lemma 1.6]. The Lemma also implies that $-(K_Y + \Gamma)$ is ample and

$$-(K_X + \Delta) = -\varphi^*(K_Y + \Gamma) + E,$$

where $E \geq 0$ is φ -exceptional, and φ^* is the pullback by birational contractions. Thus, it follows that the discrepancies of (Y, Γ) along the components of φ -divisors are at most 0. Since (Y, Γ) is klt, by [BCHM, Corollary 1.4.3] applied to

$$\mathfrak{E} = \{F \mid F \text{ is a } \varphi\text{-exceptional divisors on } X\}$$

there exists a birational morphism $f : Y' \rightarrow Y$ satisfying:

- (1) Y' is a \mathbb{Q} -factorial normal variety,
- (2) Y' and X are isomorphic in codimension one, and
- (3) there exists a \mathbb{Q} -divisor Γ' such that $f^*(K_Y + \Gamma) = K_{Y'} + \Gamma'$ and (Y', Γ') is klt.

In particular Y' is of Fano type. Thus, there exists a big \mathbb{Q} -divisor $\Theta \geq 0$ such that $(Y', \Gamma' + \Theta)$ is klt and $K_{Y'} + \Gamma' + \Theta \sim_{\mathbb{Q}} 0$.

Let Θ' be the strict transform of Θ in X . Then $(X, \Delta + \Theta')$ is klt, Θ' is big and $K_X + \Delta + \Theta' \sim_{\mathbb{R}} 0$. Thus, it follows easily from (2) that there exists a \mathbb{Q} -divisor Δ' such that $(X, \Delta + \Delta')$ is a log Fano pair (e.g. see [B, Lemma 2.4]). \square

As an immediate consequence of the previous Theorem, we obtain:

Corollary 3.2. *Let X be a \mathbb{Q} -Gorenstein normal variety.*

Then X is of Fano type if and only if $-K_X$ is big, $R(-K_X)$ is finitely generated, and $\text{Proj } R(-K_X)$ has log terminal singularities.

Remark 3.3. *Recall that if X is a normal variety, Δ is an effective \mathbb{Q} -divisor such that $L = r(K_X + \Delta)$ is ample for some $r \in \mathbb{Q}$ and C is the cone defined by L , then (X, Δ) is log Fano (in particular $r < 0$) if and only if (C, Γ) is klt, where Γ is the corresponding \mathbb{Q} -divisor on C (e.g. see [Ko, Lemma 3.1]).*

An analogue result as Theorem 3.1 holds for log Calabi-Yau pairs:

Theorem 3.4. *Let (X, Δ) be a projective log pair over an algebraic closed field k of characteristic zero. Assume that $R(X, -m(K_X + \Delta))$ is a finitely generated k -algebra for some $m \in \mathbb{N}$ and that the map*

$$\varphi : X \dashrightarrow Y := \text{Proj } R(X, -m(K_X + \Delta))$$

is a birational contraction.

Then, there exists a \mathbb{Q} -divisor $\Delta' \geq 0$ such that $(X, \Delta + \Delta')$ is a log Calabi-Yau pair if and only if the pair $(Y, \Gamma = \varphi_ \Delta)$ has only log canonical singularities.*

We begin with the following lemma:

Lemma 3.5. *Let $\mathbb{K} = \mathbb{R}$ or \mathbb{Q} . Let (X, Δ) be a log Calabi–Yau pair, for some \mathbb{K} -divisor $\Delta \geq 0$ and let X' be a normal projective variety. Suppose that there exists $\varphi : X \dashrightarrow X'$ such that φ is a birational contraction. Then $K_{X'} + \varphi_*\Delta$ is \mathbb{K} -Cartier and $(X', \varphi_*\Delta)$ is a log Calabi–Yau pair.*

Proof. Take a log resolution $f : W \rightarrow X$ of (X, Δ) which resolves the singularities of φ and let $g : W \rightarrow X'$ be the induced morphism. Let Γ be a \mathbb{K} -divisor on W such that $K_W + \Gamma = f^*(K_X + \Delta) \sim_{\mathbb{K}} 0$. Note that $g_*\Gamma = \varphi_*\Delta$. Thus $K_{X'} + \varphi_*\Delta$ is \mathbb{K} -Cartier and \mathbb{K} -linearly trivial. Let Γ' be the \mathbb{K} -divisor on W such that $K_W + \Gamma' = g^*(K_{X'} + \varphi_*\Delta) \sim_{\mathbb{K}} 0$. Thus, by the negativity lemma it follows that $\Gamma = \Gamma'$ and in particular $(X', \varphi_*\Delta)$ is log canonical. \square

Proof of Theorem 3.4. The "only if" part is an immediate consequence of the above lemma. Thus, we show the "if" part. As in the proof of Theorem 3.1, we have that $-(K_Y + \Gamma)$ is ample and

$$-(K_X + \Delta) = -\varphi^*(K_Y + \Gamma) + E,$$

where $E \geq 0$ is a φ -exceptional \mathbb{Q} -divisor. Note that the discrepancies of (Y, Γ) with respect to any φ -exceptional prime divisor is at most 0. Let $\alpha : W \rightarrow Y$ be a log resolution of (Y, Γ) which resolves the indeterminacy of φ and let $\beta : W \rightarrow X$ be the induced morphism. By assumption (Y, Γ) is log canonical. We write $\alpha^*(K_Y + \Gamma) = K_W + \Omega$, where the coefficient of Ω are less than or equal to 1. Let \mathcal{E} be the set consisting of all the α -exceptional divisors on W which are not β -exceptional and all the components of Ω with coefficient equal to one. By [F3, Theorem 4.1] applied to \mathcal{E} , there exists a birational morphism $f : Y' \rightarrow Y$ satisfying:

- (1) Y' is a \mathbb{Q} -factorial normal variety,
- (2) $Y' \dashrightarrow X$ is a birational contraction, and
- (3) there exists a \mathbb{Q} -divisor Γ' such that $g^*(K_Y + \Gamma) = (K_{Y'} + \Gamma')$ and (Y', Γ') is log canonical and $-(K_{Y'} + \Gamma')$ is semi-ample.

In particular Y' is of Calabi–Yau type. Thus, Lemma 3.5 implies that there exists $\Delta' \geq 0$ such that $(X, \Delta + \Delta')$ is a log Calabi–Yau pair. \square

Similarly to Corollary 3.2, we immediately obtain:

Corollary 3.6. *Let X be a \mathbb{Q} -Gorenstein normal variety such that $-K_X$ is big and $R(-K_X)$ is finitely generated. Then X is of Calabi–Yau type if and only if $\text{Proj } R(-K_X)$ has log canonical singularities.*

In view of Corollary 3.2, it is tempting to ask if for any variety X of Calabi–Yau type and such that $-K_X$ is big, we have that $R(X, -K_X)$ is finitely generated. However, the example below shows that this is not true in general:

Example 3.7. Let S be the blow up of \mathbb{P}^2 at 9 very general points and let $E = -K_S$. Let L_1 be an ample divisor and let $L_2 = E - L_1$. Define $X = \mathbb{P}(\mathcal{O}_S(L_1) \oplus \mathcal{O}_S(L_2))$. Then Proposition 2.10 implies that X is of Calabi–Yau type. On the other hand, we have that $\mathcal{O}_X(-K_X) = \mathcal{O}_X(2)$ is not semi-ample since E is not semi-ample. Thus, since $-K_X$ is big, it follows that $R(X, -K_X) = R(X, \mathcal{O}_X(2))$ is not finitely generated.

4. On log canonical Fano varieties

Considering Example 3.7, it seems worth studying under which condition a projective variety X is a Mori dream space. By [BCHM, Corollary 1.3.2], it follows that any variety of Fano type is a Mori dream space. The aim of this section is to study possible generalisation of this result.

We begin with the following conjecture:

Conjecture 4.1. *Let (X, Δ) be a \mathbb{Q} -factorial log canonical pair such that $-(K_X + \Delta)$ is ample. Then X is a Mori dream space.*

Note that by [BCHM, Corollary 1.3.2], the conjecture holds to be true under the assumption that (X, Δ) is dlt. We now show that the conjecture follows from the standard conjectures of the minimal model program:

Theorem 4.2. *Assume the existence of minimal models and the abundance conjecture for log canonical pairs in dimension n .*

Then Conjecture 4.1 is true in dimension n .

Proof. First, note that Kodaira vanishing for log canonical pairs (e.g. see [A] and [F2, Theorem 2.42]) implies that $H^1(X, \mathcal{O}_X) = 0$. Thus $\text{Pic}_{\mathbb{Q}}(X) \simeq N_{\mathbb{Q}}^1(X)$ (eg. see [K05, Corollary 9.5.13]). Let L_1, \dots, L_{ℓ} be a basis of $\text{Pic}_{\mathbb{Q}}(X)$. Then, for any $i = 1, \dots, \ell$, after possibly rescaling L_i , there exist effective \mathbb{Q} -divisors Δ_i such that (X, Δ_i) is log canonical and $L_i \sim_{\mathbb{Q}} K_X + \Delta_i$.

We claim that the ring

$$\bigoplus_{(m_1, \dots, m_{\ell}) \in \mathbb{N}} H^0(X, \sum_{i=1}^{\ell} m_i m (K_X + \Delta_i))$$

is finitely generated for any positive integer m such that $m(K_X + \Delta_i)$ is Cartier for any $i = 1, \dots, \ell$. Note that the claim implies that X is a Mori dream space by [HK, Theorem 2.13].

We now proceed with the proof of the claim. After taking a log resolution, we may assume that (X, Δ_i) is log smooth for any $i = 1, \dots, \ell$. Let $V \subseteq \text{Div}_{\mathbb{R}}(X)$ be the subspace spanned by the components of $\Delta_1, \dots, \Delta_{\ell}$. By [SC, Theorem 3.4] or [Ka, Theorem 3], there exist $\varphi_j: X \dashrightarrow Y_j$, with $j = 1, \dots, p$ such that if

$\Delta \in V$ is such that (X, Δ) is log canonical and $K_X + \Delta$ is pseudo-effective, then φ_j is the log terminal model of (X, Δ) for some $j = 1, \dots, p$. Thus, we obtain the claim, by using the same methods as in the proof of [BCHM, Corollary 1.1.9]. \square

In particular, we obtain:

Corollary 4.3. *Let (X, Δ) a 3-dimensional log canonical pair such that $-(K_X + \Delta)$ is ample. Then X is a Mori dream space.*

Note that, in general, a log canonical weak Fano pair is not necessarily a Mori dream space (see [G, Section 5]). The following example shows that even assuming that (X, Δ) is a log canonical pair such that $-(K_X + \Delta)$ is semi-ample and big, then X is not necessarily a Mori dream space, even assuming $H^1(X, \mathcal{O}_X) = 0$.

Example 4.4. Let S be a K3 surface which is not a Mori dream space. Let A be a very ample divisor on S and let $X = \mathbb{P}(\mathcal{O}_S \oplus \mathcal{O}_S(A))$. Thus, [Ow, Theorem 1.1] implies that X is not a Mori dream space. On the other hand, it is easy to check that if E is the section corresponding to the summand \mathcal{O}_S then (X, E) is log canonical and $-(K_X + E)$ is semi-ample and big. Furthermore, we have $H^1(X, \mathcal{O}_X) = 0$.

5. On varieties of Fano type

The purpose to this section is to explore possible characterizations of varieties of Fano type. We show two examples which show that some of the questions raised before about this problem have a negative answer.

Example 5.1. Assume that (X, Δ) is a klt pair such that $-(K_X + \Delta)$ is movable and big. Then X is not necessarily of Fano type.

Indeed, let S be a smooth projective variety of general type such that $H^1(S, \mathcal{O}_S) = 0$ and such that the Picard number of S is one. Let L_1 and L_2 be ample divisors on S , and let

$$\mathcal{E} = \mathcal{O}_S(L_1) \oplus \mathcal{O}_S(L_2) \oplus \mathcal{O}_S(-K_S - L_1 - L_2).$$

Then the anti-canonical divisor of $X := \mathbb{P}_S(\mathcal{E})$ is movable and big but since X is not rationally connected, it follows that X is not of Fano type. Note that Proposition 2.6 implies that X is a Mori dream space.

The following question is motivated by the birational Borisov-Alexeev-Borison conjecture. In [Od], Okada shows that the class of rationally connected varieties and of Fano type are birationally unbounded. Thus, it is natural to ask if these classes are the same up to birational equivalence. Recall that any variety of Fano type is rationally connected (see [Zh] and [HM]) and all smooth rationally

connected surface are rational.

Question 5.2. *Let X be a rationally connected variety. Then is X birationally equivalent to a variety of Fano type?*

Acknowledgements

The second author discussed several problems related to the examples in Section 5 with Frédéric Campana, Hiromichi Takagi, Takuzo Okada and Kiwamu Watanabe while he was a Ph.D student. Some of the questions considered in this paper were raised during an AIM workshop in May 2012. We would also like to thank Ivan Cheltsov, Anne-Sophie Kaloghiros, Ana-Maria Castravet, James M^cKernan, John Christian Ottem and Vyacheslav Vladimirovich Shokurov for several useful discussions. We are particularly thankful to Ana-Maria Castravet for pointing out an error in a previous version of this notes. We also would like to thank DongSeon Hwang for informing us that Theorem 1.1 was already known in dimension two [HP]. Finally, we would like to thank the referee for many useful comments.

References

- [A] F. Ambro, *Quasi-log varieties*, Tr. Mat. Inst. Steklova **240** (2003), Biratsion. Geom. Linein. Sist. Konechno Porozhdennye Algebr, 220–239; translation in Proc. Steklov Inst. Math., no.1, **240**, 214–233 (2003).
- [B] C. Birkar, *Singularities on the base of a Fano type fibration*, preprint arXiv:1210.2658 (2012).
- [BCHM] C. Birkar, P. Cascini, C. D. Hacon and J. M^cKernan, *Existence of minimal models for varieties of log general type*, J. Amer. Math. Soc. **23**, 405–468 (2010).
- [Br] M. Brown, *Singularities of Cox rings of Fano varieties*, J. Math. Pures Appl., <http://dx.doi.org/10.1016/j.matpur.2012.10.003> (2012).
- [Ka] Y. Kawamata, *Remarks on the cone of divisors. Classification of algebraic varieties*, EMS Ser. Congr. Rep., Eur. Math. Soc., Zürich, 317–325 (2011).
- [KaMM] Y. Kawamata, K. Matsuda and K. Matsuki, *Introduction to the minimal model problem*, Algebraic geometry, Sendai, Adv. Stud. Pure Math., 10, North-Holland, Amsterdam, 283–360 (1987).
- [Ko] J. Kollár, *Singularities of the Minimal Model Program*, Princeton University, New Jersey (2013).
- [KoM] J. Kollár and S. Mori, *Birational geometry of algebraic varieties*, Cambridge Tracts in Math., **134** (1998).
- [KKL] A. Kuronya, A.-S. Kaloghiros, and V. Lazić, *Finite generation and geography of models*, to appear in Minimal models and extremal rays, Advanced Studies in Pure Mathematics, Mathematical Society of Japan, Tokyo.
- [K05] S. Kleiman, *The Picard scheme*, Fundamental Algebraic Geometry: Grothendieck’s FGA Explained, Mathematical Surveys and Monographs, **123** (2005).
- [F1] O. Fujino, *Fundamental theorems for the log minimal model program*, Publ. Res. Inst. Math. Sci. **47**, no.3, 727–789 (2011).

- [F2] O. Fujino, *Introduction to the log minimal model program for log canonical pairs*, preprint, arXiv:0907.1506.
- [F3] O. Fujino, *Semi-stable minimal model program for varieties with trivial canonical divisor*. Proc. Japan Acad. Ser. A Math. Sci. **87**, no.3, 25–30 (2011).
- [G] Y. Gongyo, *On weak Fano varieties with log canonical singularities*, J. Reine Angew. Math. **665**, 237–252 (2012).
- [GOST] Y. Gongyo, S. Okawa, A. Sannai, and S. Takagi, *Characterization of varieties of Fano type via singularities of Cox rings*, (2011), to appear in JAG.
- [HM] C. D. Hacon and J. McKernan, *On Shokurov’s rational connectedness conjecture*, Duke Math. J. **138**, no. 1, 119–136 (2000).
- [HK] Y. Hu and S. Keel, *Mori Dream Spaces and GIT*, Michigan Math. J. **48**, 331–348 (2000).
- [HP] D-S. Hwang and J. Park, *Characterization of log del pezzo pairs via anticanonical models*, Preprint (2013).
- [M] J. McKernan, *Mori dream spaces*, Jpn. J. Math. **5** no.1, 127–151 (2010).
- [N] N. Nakayama, *Zariski decomposition and abundance*, MSJ Memoirs, **14**. Mathematical Society of Japan, Tokyo, (2004).
- [Od] T. Okada, *On the birational unboundedness of higher dimensional Q -Fano varieties*, Math. Ann. **345**, no.1, 195–212 (2009).
- [Ow] S. Okawa, *On images of Mori dream spaces*, preprint, arXiv:1104.1326 (2011).
- [PS] Yu. G. Prokhorov, V. V. Shokurov, *Towards the second main theorem on complements*, J. Algebraic Geom. **18**, no.1, 151–199 (2009).
- [Sk1] F. Sakai, *D -dimensions of algebraic surfaces and numerically effective divisors*, Compositio Math. **48** no.1, 101–118 (1983).
- [Sk2] F. Sakai, *Anticanonical models of rational surfaces*, Math. Ann. **269**, no.3, 389–410 (1983).
- [Sk3] F. Sakai, *Anti-Kodaira dimension of ruled surfaces*, Sci. Rep. Saitama Univ. Ser. A **10**, no.2, 1–7. (1982)
- [Sh] V. V. Shokurov, *3-fold log models*, J. Math. Sci. **81**, no.3, 2667–2699 (1996).
- [SC] V. V. Shokurov and S. R. Choi, *Geography of log models: theory and applications*, Cent. Eur. J. Math. **9**, no.3, 489–534 (2011).
- [TVAV] D. Testa, A. Várilly-Alvarado, M. Velasco, *Big rational surfaces*, Math. Ann. **351**, no.1, 95–107 (2011).
- [Zh] Q. Zhang, *Rational connectedness of log Q -Fano varieties*, J. Reine Angew. Math. **590**, 131–142 (2006).

Paolo Cascini
 Department of Mathematics,
 Imperial College London, 180 Queen’s Gate, London SW7 2AZ, UK
 e-mail: p.cascini@imperial.ac.uk

Yoshinori Gongyo
 Graduate School of Mathematical Sciences,
 the University of Tokyo,
 3–8–1 Komaba, Meguro-ku, Tokyo 153–8914, Japan.
 e-mail: gongyo@ms.u-tokyo.ac.jp

Department of Mathematics,
 Imperial College London, 180 Queen’s Gate, London SW7 2AZ, UK
 e-mail: y.gongyo@imperial.ac.uk